

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
14 October 2004 (14.10.2004)

PCT

(10) International Publication Number
WO 2004/087658 A1

(51) International Patent Classification⁷: **C07D 209/34**,
A61K 31/405, C07D 209/54, 209/08, A61P 25/08

(21) International Application Number:
PCT/EP2004/002691

(22) International Filing Date: 16 March 2004 (16.03.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
03007214.4 31 March 2003 (31.03.2003) EP

(71) Applicant (for all designated States except US): **UCB, S.A.**
[BE/BE]; Allée de la Recherche, 60, B-1070 Brussels (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **STARCK, Jean-Philippe** [FR/FR]; 2, rue des Tilleuls, F-67270 Gougenheim (FR). **KENDA, Benoît** [BE/BE]; Trieux des Frênes 5, B-5080 Emynes (BE).

(74) Agents: **LECHIEN, Monique** et al.; UCB, S.A., Allée de la Recherche, 60, B-1070 Brussels (BE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- with amended claims

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: INDOLONE-ACETAMIDE DERIVATIVES, PROCESSES FOR PREPARING THEM AND THEIR USES

(57) Abstract: The present invention relates to indolone-acetamide derivatives, processes for preparing them, pharmaceutical compositions containing them and their use as for the treatment of epilepsy, epileptogenesis, seizure disorders and convulsion.



WO 2004/087658 A1

INDOLONE-ACETAMIDE DERIVATIVES, PROCESSES FOR PREPARING THEM AND THEIR USES

The present invention concerns indolone-acetamide derivatives, processes for preparing them, pharmaceutical compositions containing them and their use as pharmaceuticals.


European Patent No. 0 162 036 B1 discloses the compound (S)- α -ethyl-2-oxo-1-pyrrolidine acetamide, which is known under the International Nonproprietary Name of levetiracetam.

Levetiracetam, a laevorotatory compound, is disclosed as a protective agent for the treatment and prevention of hypoxic and ischemic type aggressions of the central nervous system. This compound is also effective in the treatment of epilepsy, a therapeutic indication for which it has been demonstrated that its dextrorotatory enantiomer (R)- α -ethyl-2-oxo-1-pyrrolidine acetamide, also known from European Patent No. 0 165 919 B1, completely lacks activity (A.J. GOWER *et al.*, Eur. J. Pharmacol., 222, (1992), 193-203).

Russian patent application SU 841264 discloses 2-(2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide and its anticonvulsant activity.

It has now surprisingly been found that certain indolone-acetamide derivatives demonstrate markedly improved therapeutic properties.

In one aspect the invention therefore provides a compound having the formula I or a pharmaceutically acceptable salt thereof or stereoisomeric forms thereof,




wherein

R¹ is hydrogen,


R² is hydrogen or C1-20-alkyl,

R³ is hydrogen, C1-20-alkyl, C4-8-cycloalkyl, C5-8-cycloalkenyl, aryl, aromatic or non aromatic heterocycle, C1-20-alkoxy, or a group of formula -W-R⁸,

R^{3a} is hydrogen, C1-20-alkyl or a group of formula:



or NR^3R^{3a} is a group of formula



R^4 is hydrogen,

R^5 is hydrogen; nitro; halogen; azido; cyano; $-\text{S}-\text{C1-4-alkyl}$; $-\text{SO}-\text{C1-4-alkyl}$;

5 $-\text{SO}_2-\text{C1-4-alkyl}$; $-\text{SONH}_2$; C1-20-alkyl unsubstituted or substituted by halogen; or C1-20-alkoxy unsubstituted or substituted by halogen,

R^6 is hydrogen, C1-20-alkyl or halogen,

R^7 is hydrogen, C1-20-alkyl or halogen,

W is C1-12-alkylene, $-\text{NH}-$ or $-\text{NHC}(=\text{O})-$,

10 X is O, S or NH,

Y is O, S, $-\text{CR}^{12}\text{R}^{13}-$, $-\text{NR}^{14}-$ or $-\text{C}(=\text{O})-$,

R^8 is aryl or heterocycle,

R^9 , R^{10} , R^{10a} and R^{11} are independently selected from hydrogen, C1-4-alkyl, halogen, hydroxy or methoxycarbonyl,

15 or R^{10} and R^{10a} together form a C3-6-alkylene,

R^{12} is hydrogen, C1-4-alkyl, halogen or hydroxy,

R^{13} is hydrogen,

or $\text{CR}^{12}\text{R}^{13}$ is dioxolanyl,

R^{14} is aryl, heterocycle or a group of formula $-\text{V}-\text{R}^{15}$,

20 V is C1-12-alkylene,

R^{15} is aryl or heterocycle,


m is 1 to 4,

n is 0 or 1,

and at least one of R^5 , R^6 or R^7 is different from hydrogen when R^2 is hydrogen, R^3 is

25 H or 2,6-diisopropylphenyl, and R^{3a} is H.

In another aspect the invention provides a compound having the formula I or a pharmaceutically acceptable salt thereof or stereoisomeric forms thereof,




30 wherein

R¹ is hydrogen,


R² is hydrogen or C1-20-alkyl,

R³ is hydrogen, C1-20-alkyl, C4-8-cycloalkyl, C5-8-cycloalkenyl, aryl, aromatic or non aromatic heterocycle, C1-20-alkoxy, or a group of formula -W-R⁸,

5 R^{3a} is hydrogen, C1-20-alkyl or a group of formula:



or NR³R^{3a} is a group of formula



R⁴ is hydrogen,

10 R⁵ is hydrogen; nitro; halogen; C1-20-alkyl unsubstituted or substituted by halogen; or C1-20-alkoxy unsubstituted or substituted by halogen,

R⁶ is hydrogen, C1-20-alkyl or halogen,

R⁷ is hydrogen, C1-20-alkyl or halogen,

W is C1-12-alkylene, -NH- or -NHC(=O)-,

15 X is O, S or NH,

Y is O, S, -CR¹²R¹³-, -NR¹⁴- or -C(=O)-,

R⁸ is aryl or heterocycle,

R⁹, R¹⁰, R^{10a} and R¹¹ are independently selected from hydrogen, C1-4-alkyl, halogen, hydroxy or methoxycarbonyl,

20 or R¹⁰ and R^{10a} together form a C3-6-alkylene,

R¹² is hydrogen, C1-4-alkyl, halogen or hydroxy,

R¹³ is hydrogen,

or CR¹²R¹³ is dioxolanyl,

R¹⁴ is aryl, heterocycle or a group of formula -V-R¹⁵,

25 V is C1-12-alkylene,

R¹⁵ is aryl or heterocycle,

m is 1 to 4,

n is 0 or 1,

and at least one of R⁵, R⁶ or R⁷ is different from hydrogen when R² is hydrogen, R³ is

30 H or 2,6-diisopropylphenyl, and R^{3a} is H.

The term "alkyl", as used herein, is defined as including saturated, monovalent hydrocarbon radicals having straight, branched or cyclic moieties or combinations thereof and containing 1-20 carbon atoms, preferably 1-6 carbon atoms and more preferably 1-4 carbon atoms for non-cyclic alkyl and 3-8 carbon atoms for cycloalkyl.

5 Alkyl moieties may optionally be substituted by 1 to 5 substituents independently selected from halogen, hydroxy, alkoxy, alkoxycarbonyl, ester or alkylamino. Preferred alkyl groups are methyl, ethyl, n-propyl, isopropyl, trifluoromethyl, n-butyl, 2-fluoroethyl, 3-hydroxypropyl, 3-hydroxy-2,2-dimethylpropyl, 1-(hydroxymethyl)propyl, 3,3,3-trifluoro-2-hydroxypropyl, 3-ethoxypropyl, 2-ethoxy-2-oxoethyl and 3-
10 (dimethylamino)propyl.

The term "cycloalkyl", as used herein, refers to a monovalent group of 3 to 18 carbon atoms, preferably 4-8 carbon atoms, derived from a saturated cyclic or polycyclic hydrocarbon which may be substituted by any suitable group including but not limited to one or more moieties selected from groups as described above for the
15 alkyl groups. Preferred cycloalkyl group is cycloheptyl.

The term "alkylene", as used herein, represents a divalent alkyl group, having straight or branched moieties, containing 1-12 carbon atoms, preferably 1-6 carbon atoms, and being optionally substituted with any suitable group, including but not limited to one or more moieties selected from groups as described above for the alkyl
20 groups. Preferred alkylene groups are methylene, ethylene, hydroxyethylene, trimethylene or propylene.

The term "cycloalkenyl", as used herein, is defined as a cyclic unsaturated hydrocarbon radical having at least one double bond, containing 4-20 carbon atoms, preferably 5-8 carbon atoms, and being optionally substituted with any suitable
25 group, including but not limited to one or more moieties selected from groups as described above for the alkyl groups. Preferred cycloalkenyl group is 6-(hydroxymethyl)cyclohex-3-en-1-yl.

The term "aryl", as used herein, is defined as including an organic radical derived from an aromatic hydrocarbon consisting of 1-3 rings and containing 6-30
30 carbon atoms by removal of one hydrogen, such as phenyl and naphthyl each optionally substituted by 1 to 5 substituents independently selected from halogen, hydroxy, nitro, C1-6-alkyl, C1-6-alkoxy, C1-6-alkylsulfonyl, trifluoromethylthio or pyridinylalkyl. Aryl radicals are preferably phenyl radicals. Preferred aryl groups are phenyl, 3-hydroxyphenyl, 3-fluorophenyl, 3-methylphenyl, 4-methylphenyl, 4-
35 hydroxyphenyl, 4-hydroxy-3-methoxyphenyl, 3-(2-pyridin-2-ylethyl)phenyl, 3,4-dimethylphenyl, 4-tert-butylphenyl, 4-methylsulfonylphenyl, 2-nitrophenyl, 2-chloro-6-fluorophenyl, 2-[(trifluoromethyl)thio]phenyl, 2-chlorophenyl or 4-bromophenyl.

The term "halogen", as used herein, includes an atom of Cl, Br, F, I.

The term "nitro", as used herein, represents a group of the formula -NO₂.

The term "hydroxy", as used herein, represents a group of the formula -OH.

5 The term "alkoxy", as used herein, represents a group of formula -OR^b wherein R^b is an alkyl group, as defined above.

The term "ester", as used herein, represents a group of formula -COOR^c wherein R^c is an alkyl group or an aryl group, as defined above.

The term "alkoxycarbonyl", as used herein, represents a group of formula -COOR^d wherein R^d is an alkyl group, as defined above.

10 The term "amino", as used herein, represents a group of the formula -NH₂.

The term "alkylamino", as used herein, represents a group of formula -NHR^e or -NR^eR^f wherein R^e and R^f are alkyl group as defined above.

The term alkylsulfonyl, as used herein is defined as representing a group of formula -SO₂-R^g, wherein R^g is C1-4-alkyl.

15 The term "heterocycle", as used herein is defined as including an aromatic or non aromatic cycloalkyl or cycloalkenyl moiety as defined above, having at least one O, S and/or N atom interrupting the carbocyclic ring structure and optionally, one of the carbon of the carbocyclic ring structure may be replaced by a carbonyl.

Non-limiting examples of aromatic heterocycles are pyrazolyl, furyl, imidazolyl, triazolyl, oxazolyl, pyridinyl, pyrrolyl, thienyl, isothiazolyl, benzimidazolyl, tetrazolyl, isooxazolyl, oxazolyl, thiazolyl, 1,2,4-thiadiazolyl, oxadiazole, pyridazinyl, pyrimidinyl, pyrazinyl, isoindolyl, triazolopyridinyl, imidazolopyridinyl, pyrrolopyrimidinyl, pyrazolopyrimidinyl, quinazolinyl, quinoliziny, naphthyridinyl, quinolyl, isoquinolyl, isobenzofuranyl, benzothienyl, indolyl, indoliziny, purinyl, carbazolyl, thieno(2,3-
25 b)furanyl, thianthrenyl, benzothiazolyl, benzoxazolyl, cinnoliny, quinoxaliny, phenothiaziny, isochromanyl and xanthenyl, optionally substituted by 1 to 5 substituents independently selected from halogen, hydroxy, thiol, amino, nitro, cyano, azido, C1-6-alkoxy, C1-6-alkylthio, C1-6-alkyl, C1-6-haloalkyl, formyl or ester. More preferred aromatic heterocycles are pyrazolyl, furyl, imidazolyl, triazolyl, oxazolyl and
30 pyridinyl.

Non-limiting examples of non aromatic heterocycles are tetrahydrofuranyl, piperidinyl, piperidyl, piperazinyl, imidazolidinyl, morpholiny, thiomorpholiny, pyrrolidinyl, thiazolidinyl, indoliny, tetrahydrobenzazociny, dihydroisochromenyl, tetrahydropyranyl, oxooctahydroquinoliny, dioxolanyl, 1-oxaspiro(4.5)dec-2-yl, pyrrolidinyl, 2-oxo-pyrrolidinyl, 8-thiabicyclo[3.2.1]cyclooctanyl, 1,4-dithiepanyl, tetrahydro-2H-thiopyranyl, azepanyl and azocanyl, optionally substituted by 1 to 5
35 substituents independently selected from halogen, hydroxy, thiol, amino, nitro, cyano,

azido, C1-6-alkoxy, C1-6-alkylthio, C1-6-alkyl, C1-6-haloalkyl, formyl or ester. More preferred non aromatic heterocycles are tetrahydrofuran-1-yl, piperidin-1-yl, piperidin-2-yl, piperazin-1-yl, imidazolidin-1-yl, morpholin-1-yl, thiomorpholin-1-yl, pyrrolidin-1-yl, thiazolidin-1-yl, indolin-1-yl, tetrahydro-1-benzazocin-1(2H)-yl, 3,4-dihydro-1H-isochromen-1-yl, tetrahydropyranyl, oxooctahydroquinolin-1-yl and dioxolanyl. The term "heterocycle" also includes bicyclic, tricyclic and tetracyclic, spiro groups in which any of the above heterocyclic rings is fused to one or two rings independently selected from an aryl ring, a cycloalkyl ring, a cycloalkenyl ring or another monocyclic heterocyclic ring or where a monocyclic heterocyclic group is bridged by an alkylene group, such as quinuclidinyl, 7-azabicyclo(2.2.1)heptan-1-yl, 7-oxabicyclo(2.2.1)heptan-1-yl and 8-azabicyclo(3.2.1)octan-1-yl.

The term "pyridinylalkyl", as used herein, represents a group of formula $-R^h$ -pyridinyl in which R^h is C1-4-alkylene.

The term "azido" as used herein, represents a group of the formula $-N_3$.

The term "cyano" as used herein, represents a group of the formula $-CN$.

Generally, R^2 is hydrogen or C1-4-alkyl.

Preferably, R^2 is hydrogen, methyl or ethyl. More preferably, R^2 is hydrogen or methyl.

Generally, R^3 is hydrogen; C1-6-alkyl unsubstituted or substituted by 1 to 5 substituents selected from halogen, hydroxy, alkoxy, alkoxycarbonyl or alkylamino; C5-7-cycloalkyl; (hydroxymethyl)cyclohexenyl; phenyl unsubstituted or substituted by 1 to 5 substituents selected from halogen, C1-4-alkyl, hydroxy, methoxy, nitro, methylsulfonyl, trifluoromethylthio or pyridinylalkyl; pyridinyl unsubstituted or substituted by methoxy; triazolyl; C1-4-alkoxy; or a group of formula $-W-R^8$ wherein:


Generally, W is C1-4-alkylene unsubstituted or substituted by halogen, hydroxy, C1-4-alkyl or alkoxy; $-NH-$; or $-NHC(=O)-$; and

R^8 is phenyl unsubstituted or substituted by 1 to 5 substituents selected from halogen, C1-4-alkyl, hydroxy, methoxy, nitro, methylsulfonyl or trifluoromethylthio; furyl unsubstituted or substituted by methyl; pyrazolyl; pyridinyl; morpholinyl; tetrahydrobenzazocinyl; piperidinyl unsubstituted or substituted by methyl; dihydroisochromenyl or dihydroimidazolyl.

Preferably, R^3 is hydrogen, n-butyl, cycloheptyl, 2-fluoroethyl, 3-hydroxypropyl, 3-hydroxy-2,2-dimethylpropyl, 1-(hydroxymethyl)propyl, 3,3,3-trifluoro-2-hydroxypropyl, 3-ethoxypropyl, 2-ethoxy-2-oxoethyl, 3-(dimethylamino)propyl, 6-(hydroxymethyl)cyclohex-3-en-1-yl, 3-hydroxyphenyl, 3-fluorophenyl, 3-(2-pyridin-2-ylethyl)phenyl, 3,4-dimethylphenyl, 4-tert-butylphenyl, benzyl, 4-hydroxy-3-methoxybenzyl, 4-methylsulfonylbenzyl, 2-nitrobenzyl, 2-chloro-

6-fluorobenzyl, 2-[(trifluoromethyl)thio]benzyl, 2-hydroxy-2-phenylethyl, 2-(3,4-dimethoxyphenyl)ethyl, 2-(2-chlorophenyl)ethyl, 2-(4-methylphenyl)ethyl, (4-bromophenyl)amino, pyridin-3-yl, 6-methoxypyridin-3-yl, 4H-1,2,4-triazol-3-yl, pyridin-4-ylmethyl, (5-methyl-2-furyl)methyl, 3-(1H-pyrazol-1-yl)propyl, 2-morpholin-4-ylethyl, 2-((3,4,5,6-tetrahydro-1-benzazocin-1(2H)-yl)propyl, 2-(2-methylpiperidin-1-yl)ethyl, 3,4-dihydro-1H-isochromen-1-ylmethyl, methoxy, (4-pyridinylcarbonyl)amino or 4,5-dihydro-1H-imidazol-2-ylamino. More preferably, R³ is hydrogen.


Generally, R^{3a} is hydrogen, C1-4-alkyl or a group of formula



wherein m is 1 to 4.

Preferably, R^{3a} is hydrogen, methyl or tetrahydrofuran-2-ylmethyl. More preferably, R^{3a} is hydrogen.

In another embodiment, NR³R^{3a} is piperidinyl unsubstituted or substituted by hydroxy; thiomorpholinyl; thiazolidinyl unsubstituted or substituted by C1-4-alkoxycarbonyl; 2,5-dihydro-1H-pyrrol-1-yl; 1,4-dioxo-8-azaspiro[4.5]dec-8-yl; 4-oxooctahydro-1(2H)-quinolinyl; or a group of formula



wherein R¹⁴ is pyridinyl; phenyl unsubstituted or substituted by halogen, hydroxy, C1-4-alkyl; or a group of formula -V-R¹⁵ wherein V is unsubstituted C1-4-alkylene and R¹⁵ is phenyl or morpholinyl.

In a preferred embodiment, NR³R^{3a} is 4-pyridin-2-ylpiperazin-1-yl, 4-(3-methylphenyl)piperazin-1-yl, 4-(4-hydroxyphenyl)piperazin-1-yl, 4-(2-phenylethyl)piperazin-1-yl, 4-(2-morpholin-4-ylethyl)piperazin-1-yl, 3-hydroxypiperidin-1-yl, thiomorpholin-4-yl, 4-methoxycarbonyl-1,3-thiazolidin-3-yl, 2,5-dihydro-1H-pyrrol-1-yl, 1,4-dioxo-8-azaspiro[4.5]dec-8-yl or 4-oxooctahydro-1(2H)-quinolinyl.

Generally, R⁵ is hydrogen, nitro, halogen, C1-4-alkyl, unsubstituted or substituted by halogen, or C1-4-alkoxy unsubstituted or substituted by halogen.

Preferably, R⁵ is hydrogen, methyl, ethyl, trifluoromethyl, trifluoromethoxy, n-propyl, isopropyl, nitro, or halogen. More preferably, R⁵ is halogen or trifluoromethyl.

Generally, R⁶ is hydrogen, C1-6-alkyl or halogen.


Preferably, R⁶ is hydrogen, methyl or Cl. More preferably, R⁶ is hydrogen.

Generally, R⁷ is hydrogen, methyl or halogen.

Preferably, R⁷ is hydrogen, methyl, Br, F or Cl. More preferably, R⁷ is hydrogen, Br or F.


Combinations of one or more of these preferred compound groups are especially preferred.

In a preferred embodiment, the invention provides a compound having the
5 formula I or a pharmaceutically acceptable salt thereof or stereoisomeric forms thereof,




wherein

- R¹ is hydrogen,
10 R² is hydrogen or C1-4-alkyl,
R³ is hydrogen; C1-6-alkyl unsubstituted or substituted by 1 to 5 substituents selected from halogen, hydroxy, alkoxy, alkoxycarbonyl or alkylamino; C5-7-cycloalkyl; (hydroxymethyl)cyclohexenyl; phenyl unsubstituted or substituted by 1 to 5 substituents selected from halogen, C1-4-alkyl, hydroxy, methoxy, nitro,
15 methylsulfonyl, trifluoromethylthio or pyridinylalkyl; pyridinyl unsubstituted or substituted by methoxy; triazolyl; C1-4-alkoxy; or a group of formula -W-R⁸,
R^{3a} is hydrogen, C1-4-alkyl or a group of formula



- or NR³R^{3a} is piperidinyl unsubstituted or substituted by hydroxy; thiomorpholinyl;
20 thiazolidinyl unsubstituted or substituted by C1-4-alkoxycarbonyl; 2,5-dihydro-1H-pyrrol-1-yl; 1,4-dioxo-8-azaspiro[4.5]dec-8-yl; 4-oxooctahydro-1(2H)-quinolinyl; or a group of formula



- R⁴ is hydrogen,
25 R⁵ is hydrogen; nitro; halogen; C1-4-alkyl, unsubstituted or substituted by halogen; or C1-4-alkoxy unsubstituted or substituted by halogen,
R⁶ is hydrogen, C1-6-alkyl or halogen,
R⁷ is hydrogen, methyl or halogen,
W is C1-4-alkylene unsubstituted or substituted by halogen, hydroxy, C1-4-alkyl or alkoxy; -NH-; or -NHC(=O)-,
30

R⁸ is phenyl unsubstituted or substituted by 1 to 5 substituents selected from halogen, C1-4-alkyl, hydroxy, methoxy, nitro, methylsulfonyl or trifluoromethylthio; furyl unsubstituted or substituted by methyl; pyrazolyl; pyridinyl; morpholinyl; tetrahydrobenzazocinyl; piperidinyl unsubstituted or substituted by methyl;

5 dihydroisochromenyl or dihydroimidazolyl,

R¹⁴ is pyridinyl; phenyl unsubstituted or substituted by halogen, hydroxy, C1-4-alkyl; or a group of formula -V-R¹⁵,


V is unsubstituted C1-4-alkylene,

R¹⁵ is phenyl or morpholinyl,

10 m is 1 to 4,

and at least one of R⁵, R⁶ or R⁷ is different from hydrogen when R² is hydrogen, R³ is H or 2,6-diisopropylphenyl, and R^{3a} is H.

In a more preferred embodiment, the invention provides a compound having the formula I or a pharmaceutically acceptable salt thereof or stereoisomeric forms thereof,



wherein

R¹ is hydrogen,

R² is hydrogen, methyl or ethyl,

20 R³ is hydrogen, n-butyl, cycloheptyl, 2-fluoroethyl, 3-hydroxypropyl, 3-hydroxy-2,2-dimethylpropyl, 1-(hydroxymethyl)propyl, 3,3,3-trifluoro-2-hydroxypropyl, 3-ethoxypropyl, 2-ethoxy-2-oxoethyl, 3-(dimethylamino)propyl, 6-(hydroxymethyl)cyclohex-3-en-1-yl, 3-hydroxyphenyl, 3-fluorophenyl, 3-(2-pyridin-2-ylethyl)phenyl, 3,4-dimethylphenyl, 4-tert-butylphenyl, benzyl, 4-hydroxy-3-methoxybenzyl, 4-methylsulfonylbenzyl, 2-nitrobenzyl, 2-chloro-6-fluorobenzyl, 2-
25 [(trifluoromethyl)thio]benzyl, 2-hydroxy-2-phenylethyl, 2-(3,4-dimethoxyphenyl)ethyl, 2-(2-chlorophenyl)ethyl, 2-(4-methylphenyl)ethyl, (4-bromophenyl)amino, pyridin-3-yl, 6-methoxypyridin-3-yl, 4H-1,2,4-triazol-3-yl, pyridin-4-ylmethyl, (5-methyl-2-furyl)methyl, 3-(1H-pyrazol-1-yl)propyl, 2-morpholin-4-ylethyl, 2-((3,4,5,6-tetrahydro-
30 1-benzazocin-1(2H)-yl)propyl, 2-(2-methylpiperidin-1-yl)ethyl, 3,4-dihydro-1H-isochromen-1-ylmethyl, methoxy, (4-pyridinylcarbonyl)amino or 4,5-dihydro-1H-imidazol-2-ylamino,
R^{3a} is hydrogen, methyl or tetrahydrofuran-2-ylmethyl,

or NR³R^{3a} 4-pyridin-2-ylpiperazin-1-yl, 4-(3-methylphenyl)piperazin-1-yl, 4-(4-hydroxyphenyl)piperazin-1-yl, 4-(2-phenylethyl)piperazin-1-yl, 4-(2-morpholin-4-ylethyl)piperazin-1-yl, 3-hydroxypiperidin-1-yl, thiomorpholin-4-yl, 4-methoxycarbonyl-1,3-thiazolidin-3-yl, 2,5-dihydro-1H-pyrrol-1-yl, 1,4-dioxo-8-azaspiro[4.5]dec-8-yl or 4-oxooctahydro-1(2H)-quinolinyl,
 5 R⁴ is hydrogen,
 R⁵ is hydrogen, methyl, ethyl, trifluoromethyl, trifluoromethoxy, n-propyl, isopropyl, nitro or halogen,
 R⁶ is hydrogen, methyl or Cl,
 10 R⁷ is hydrogen, methyl, Br, F or Cl,
 and at least one of R⁵, R⁶ or R⁷ is different from hydrogen when R² is hydrogen, R³ is H or 2,6-diisopropylphenyl, and R^{3a} is H.

More preferably, R² is hydrogen or methyl, R³ is hydrogen, R^{3a} is hydrogen, R⁵ is halogen or trifluoromethyl, R⁶ is hydrogen and R⁷ is hydrogen, Br or F.

15 In all the above-mentioned scopes, when R² is C1-20-alkyl, the carbon atom to which R² is attached is preferably in the "S"-configuration.

Preferred compounds are: 2-(5-iodo-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5,7-dibromo-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-nitro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-methyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; (2R)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; (2S)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; 2-[2-oxo-5-(trifluoromethoxy)-2,3-dihydro-1H-indol-1-yl]acetamide; 2-(5-isopropyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-ethyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-fluoro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5,7-dimethyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-bromo-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(2-oxo-5-propyl-2,3-dihydro-1H-indol-1-yl)acetamide; 2-[2-oxo-5-(trifluoromethyl)-2,3-dihydro-1H-indol-1-yl]acetamide; 2-(5,6-dimethyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(7-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(6-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)butanamide; (+)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)butanamide; (-)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)butanamide; 2-(5-methyl-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; (+)-2-(5-methyl-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; (-)-2-(5-methyl-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; 2-(5-bromo-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; (-)-2-(5-bromo-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; (+)-2-(5-bromo-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; 2-(5-chloro-7-fluoro-2-oxo-

- 2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3-hydroxyphenyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3-fluorophenyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[3-(2-pyridin-2-ylethyl)phenyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[6-(hydroxymethyl)cyclohex-3-en-1-yl]acetamide; 5-chloro-1-[2-oxo-2-(4-pyridin-2-ylpiperazin-1-yl)ethyl]-1,3-dihydro-2H-indol-2-one; 5-chloro-1-[2-[4-(3-methylphenyl)piperazin-1-yl]-2-oxoethyl]-1,3-dihydro-2H-indol-2-one; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(4-hydroxy-3-methoxybenzyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(pyridin-4-ylmethyl)-N-(tetrahydrofuran-2-ylmethyl)acetamide; 5-chloro-1-[2-(3-hydroxypiperidin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-indol-2-one; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N'-isonicotinoylacetohydrazide; 5-chloro-1-(2-oxo-2-thiomorpholin-4-ylethyl)-1,3-dihydro-2H-indol-2-one; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(4H-1,2,4-triazol-3-yl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[4-(methylsulfonyl)benzyl]acetamide; 1-[(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetyl]octahydroquinolin-4(1H)-one; N'-(4-bromophenyl)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetohydrazide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(6-methoxypyridin-3-yl)acetamide; N-butyl-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3-hydroxypropyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[3-(dimethylamino)propyl]acetamide; 5-chloro-1-[2-oxo-2-[4-(2-phenylethyl)piperazin-1-yl]ethyl]-1,3-dihydro-2H-indol-2-one; ethyl {[5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetyl]amino}acetate; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3-ethoxypropyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(2-fluoroethyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-methoxy-N-methylacetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3,4-dimethylphenyl)acetamide; N-(4-tert-butylphenyl)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3-hydroxy-2,2-dimethylpropyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[1-(hydroxymethyl)propyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3,3,3-trifluoro-2-hydroxypropyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(2-hydroxy-2-phenylethyl)acetamide; 5-chloro-1-[2-[4-(4-hydroxyphenyl)piperazin-1-yl]-2-oxoethyl]-1,3-dihydro-2H-indol-2-one; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(pyridin-4-ylmethyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[(5-methyl-2-furyl)methyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[3-(1H-pyrazol-1-yl)propyl]acetamide; methyl 3-[(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetyl]-1,3-thiazolidine-4-carboxylate; 5-

chloro-1-[2-(2,5-dihydro-1H-pyrrol-1-yl)-2-oxoethyl]-1,3-dihydro-2H-indol-2-one; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N'-(4,5-dihydro-1H-imidazol-2-yl)acetohydrazide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-(3,4-dimethoxyphenyl)ethyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-(2-chlorophenyl)ethyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-(4-methylphenyl)ethyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(2-morpholin-4-ylethyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-(3,4,5,6-tetrahydro-1-benzazocin-1(2H)-yl)propyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-(2-methylpiperidin-1-yl)ethyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(2-nitrobenzyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3,4-dihydro-1H-isochromen-1-ylmethyl)acetamide; N-(2-chloro-6-fluorobenzyl)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; N-benzyl-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-methylacetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-{2-[(trifluoromethyl)thio]benzyl}acetamide; 5-chloro-1-[2-(1,4-dioxo-8-azaspiro[4.5]dec-8-yl)-2-oxoethyl]-1,3-dihydro-2H-indol-2-one; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-cycloheptylacetamide; 5-chloro-1-[2-[4-(2-morpholin-4-ylethyl)piperazin-1-yl]-2-oxoethyl]-1,3-dihydro-2H-indol-2-one; and 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-pyridin-3-ylacetamide.

More preferred compounds are: 2-(5-iodo-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5,7-dibromo-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; (2S)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; 2-[2-oxo-5-(trifluoromethyl)-2,3-dihydro-1H-indol-1-yl]acetamide and 2-(5-chloro-7-fluoro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide.

In a most preferred embodiment the invention relates to a compound selected from 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide and (2S)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide.

The "pharmaceutically acceptable salts" according to the invention include therapeutically active, non-toxic base salt forms which the compounds of formula I are able to form.

The compounds of formula I containing acidic protons may be converted into their therapeutically active, non-toxic base addition salt forms, e.g. metal or amine salts, by treatment with appropriate organic and inorganic bases. Appropriate base salt forms include, for example but not limited to, ammonium salts, alkali and earth alkaline metal salts, e.g. lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. N-methyl-D-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

Conversely said salt forms can be converted into the free forms by treatment

with an appropriate acid.

Compounds of the formula I and their salts can be in the form of a solvate, which is included within the scope of the present invention. Such solvates include for example hydrates, alcoholates and the like.

5 Many of the compounds of formula I and some of their intermediates have at least one stereogenic center in their structure. This stereogenic center may be present in a R or a S configuration, said R and S notation is used in correspondence with the rules described in Pure Appl. Chem., 45 (1976) 11-30.

10 The invention also relates to all stereoisomeric forms such as enantiomeric and diastereoisomeric forms of the compounds of formula I or mixtures thereof (including all possible mixtures of stereoisomers).


Some of the compounds of formula I may also exist in tautomeric forms. Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention.

15 With respect to the present invention reference to a compound or compounds is intended to encompass that compound in each of its possible isomeric forms and mixtures thereof unless the particular isomeric form is referred to specifically.

Compounds according to the present invention may exist in different polymorphic forms. Although not explicitly indicated in the above formula, such forms
20 are intended to be included within the scope of the present invention.


The compounds of formula I according to the invention can be prepared analogously to conventional methods as understood by the person skilled in the art of synthetic organic chemistry.

25 **A.** According to one embodiment, some compounds having the general formula I may be prepared by desulfurization of a compound of formula II according to the equation:



30 This reaction may be carried out using Raney nickel in an inert solvent, preferably THF, at a temperature comprised between 0 °C and 40 °C, or as described in: Mehta L., Parrick J., Payne F., J. Chem. Research (S) (1998), 190-191.


Compounds of formula II may be prepared by alkylation of a compound of formula III with a compound of formula IV according to the equation:



wherein Hal is a halogen atom, preferably Br or Cl, and R^1 , R^2 , R^3 , R^{3a} , R^4 , R^5 and R^7 have the same definitions as described above.

This reaction may be carried out with a strong base, for example sodium hydride, at a temperature comprised between 0 and 40 °C and in an inert solvent, for example DMF under an inert atmosphere, or as described in patent GB 1,309,692 (UCB).

Compounds of formula III may be prepared by reaction of a compound of formula V with 1,2-ethanedithiol according to the equation:




wherein R^4 , R^5 and R^7 have the same definitions as described above.

This reaction may be carried out at a temperature comprised between 25 and 100 °C in an inert solvent or in acetic acid, in the presence of a Lewis acid, preferably $BF_3 \cdot Et_2O$ under an inert atmosphere.


Compounds of formula V are commercially available or may be prepared according to methods described in: Smith K., El-Hiti G.A., Hawes A.C., Synlett (1999), 945-947; Lackey K., Sternbach D.D., Synthesis (1993), 10, 993; or Organic Synthesis, Collective Volume I, Second Edition, Gilman H. & Blatt A.H., J. Wiley & Sons Inc., 327-330.

B. According to another embodiment, some compounds having the general formula I may be prepared by oxidative bromination of the corresponding indole of formula (VI) followed by the reduction of compound (VII) according to the equation:



This reaction may be carried out as described in: Marfat A., Carta M.P., Tetrahedron Lett. (1987), 28, 4027-4031.

Compounds of formula VI may be prepared by alkylation of a compound of formula VIII with a compound of formula IV according to the equation:



wherein Hal is an halogen atom, preferably Br or Cl.

This reaction may be carried out in the presence of a strong base, preferably sodium hydride, at a temperature comprised between 0 and 40 °C, in an inert solvent, for example DMF, under an inert atmosphere, or as described in patent GB 1,309,692 (UCB).


C. According to another embodiment, some compounds having the general formula I may be prepared by halogenation of the corresponding compound of formula I wherein R⁵ is a hydrogen, with a N-halosuccinimide according to the procedure described in: Castanet A.-S., Colobert F., Broutin P.-E., Tetrahedron Lett. (2002), 43, 5047-5048.

According to another embodiment, some compounds having the general formula I may be prepared analogously from the corresponding compound of formula I wherein R⁵ = R⁷ = H by using two equivalents of N-halosuccinimide.


D. According to another embodiment, some compounds having the general formula I may be prepared by nitration of the corresponding compound of formula I wherein R⁵ is a hydrogen according to procedure described in: Sun L., Rubin J.R., Kraker A.J., Showalter H.D., J. Heterocyclic Chem. (1997), 34, 1399-1405.

E. According to another embodiment, some compounds having the general formula I may be prepared by coupling of an amine of formula NHR³R^{3a} with a carboxylic acid derivative of formula IX in the presence of a coupling agent such as dicyclohexylcarbodiimide in dichloromethane or THF.

16



In one embodiment, the present invention concerns also the synthesis intermediates of formula II or stereoisomeric forms thereof,



5

wherein


R¹ is hydrogen,

R² is hydrogen or C1-20-alkyl,


R³ is hydrogen, C1-20-alkyl, C4-8-cycloalkyl, C5-8-cycloalkenyl, aryl, aromatic or non aromatic heterocycle, C1-20-alkoxy, or a group of formula -W-R⁸,

10

R^{3a} is hydrogen, C1-20-alkyl or a group of formula:



or NR³R^{3a} is a group of formula



15

R⁴ is hydrogen,

R⁵ is hydrogen; nitro; halogen; azido; cyano; -S-C1-4-alkyl; -SO-C1-4-alkyl;

-SO₂-C1-4-alkyl; -SONH₂; C1-20-alkyl unsubstituted or substituted by halogen; or C1-20-alkoxy unsubstituted or substituted by halogen,

R⁶ is hydrogen, C1-20-alkyl or halogen,

20

R⁷ is hydrogen, C1-20-alkyl or halogen,

W is C1-12-alkylene, -NH- or -NHC(=O)-,

X is O, S or NH,

Y is O, S, $-\text{CR}^{12}\text{R}^{13}-$, $-\text{NR}^{14}-$ or $-\text{C}(=\text{O})-$,

R^8 is aryl or heterocycle,

R^9 , R^{10} , R^{10a} and R^{11} are independently selected from hydrogen, C1-4-alkyl, halogen, hydroxy or methoxycarbonyl,

5 or R^{10} and R^{10a} together form a C3-6-alkylene,

R^{12} is hydrogen, C1-4-alkyl, halogen or hydroxy,

R^{13} is hydrogen,

or $\text{CR}^{12}\text{R}^{13}$ is dioxolanyl,

R^{14} is aryl, heterocycle or a group of formula $-\text{V}-\text{R}^{15}$,

10 V is C1-12-alkylene,

R^{15} is aryl or heterocycle,


m is 1 to 4,

n is 0 or 1,

and at least one of R^5 , R^6 or R^7 is different from hydrogen when R^2 is hydrogen, R^3 is

15 H or 2,6-diisopropylphenyl, and R^{3a} is H.

In another embodiment, the present invention concerns also the synthesis intermediates of formula II or stereoisomeric forms thereof,




wherein

20 R^1 is hydrogen,

R^2 is hydrogen or C1-20-alkyl,


R^3 is hydrogen, C1-20-alkyl, C4-8-cycloalkyl, C5-8-cycloalkenyl, aryl, aromatic or non aromatic heterocycle, C1-20-alkoxy, or a group of formula $-\text{W}-\text{R}^8$,

R^{3a} is hydrogen, C1-20-alkyl or a group of formula:



25

or NR^3R^{3a} is a group of formula




R^4 is hydrogen,

- R⁵ is hydrogen; nitro; halogen; C1-20-alkyl unsubstituted or substituted by halogen;
 or C1-20-alkoxy unsubstituted or substituted by halogen,
 R⁶ is hydrogen, C1-20-alkyl or halogen,
 R⁷ is hydrogen, C1-20-alkyl or halogen,
 5 W is C1-12-alkylene, -NH- or -NHC(=O)-,
 X is O, S or NH,
 Y is O, S, -CR¹²R¹³-, -NR¹⁴- or -C(=O)-,
 R⁸ is aryl or heterocycle,
 R⁹, R¹⁰, R^{10a} and R¹¹ are independently selected from hydrogen, C1-4-alkyl,
 10 halogen, hydroxy or methoxycarbonyl,
 or R¹⁰ and R^{10a} together form a C3-6-alkylene,
 R¹² is hydrogen, C1-4-alkyl, halogen or hydroxy,
 R¹³ is hydrogen,
 or CR¹²R¹³ is dioxolanyl,
 15 R¹⁴ is aryl, heterocycle or a group of formula -V-R¹⁵,
 V is C1-12-alkylene,
 R¹⁵ is aryl or heterocycle,
 m is 1 to 4,
 n is 0 or 1,
 20 and at least one of R⁵, R⁶ or R⁷ is different from hydrogen when R² is hydrogen, R³ is
 H or 2,6-diisopropylphenyl, and R^{3a} is H.

Preferably, the synthesis intermediates of formula II are selected from the
 group consisting of: 2-(5'-methyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-
 yl)acetamide; 2-[2'-oxo-5'-[(trifluoromethyl)oxy]spiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-
 25 yl]acetamide; 2-[5'-(1-methylethyl)-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-
 yl]acetamide; 2-(5'-ethyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide;
 2-(5'-fluoro-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide; 2-(5',7'-
 dimethyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide; 2-(2'-oxo-5'-
 propylspiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide; 2-[2'-oxo-5'-
 30 (trifluoromethyl)spiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl]acetamide; and 2-(5',6'-
 dimethyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide.

In one embodiment, the invention concerns also the synthesis intermediates of
 formula III or stereoisomeric forms thereof,



wherein

R⁴ is hydrogen,

R⁵ is hydrogen; nitro; azido; cyano, -S-C1-4-alkyl; -SO-C1-4-alkyl;

-SO₂-C1-4-alkyl; -SONH₂; C1-20-alkyl unsubstituted or substituted by halogen; or

5 C1-20-alkoxy unsubstituted or substituted by halogen,


R⁶ is hydrogen, C1-20-alkyl or halogen,

R⁷ is hydrogen, C1-20-alkyl or halogen,

and at least one of R⁵, R⁶ or R⁷ is different from hydrogen.

In another embodiment, the invention concerns also the synthesis

10 intermediates of formula III or stereoisomeric forms thereof,



wherein

R⁴ is hydrogen,

R⁵ is hydrogen; nitro; C1-20-alkyl unsubstituted or substituted by halogen; or C1-20-

15 alkoxy unsubstituted or substituted by halogen,

R⁶ is hydrogen, C1-20-alkyl or halogen,

R⁷ is hydrogen, C1-20-alkyl or halogen,

and at least one of R⁵, R⁶ or R⁷ is different from hydrogen.

Preferably, the synthesis intermediates of formula III are selected from the

20 group consisting of: 5'-methylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one; 5'-

[(trifluoromethyl)oxy]spiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one; 5'-(1-

methylethyl)spiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one; 5'-ethylspiro[1,3-dithiolane-

2,3'-indol]-2'(1'H)-one; 5'-fluorospiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one; 5',7'-


dimethylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one; 5'-propylspiro[1,3-dithiolane-2,3'-

25 indol]-2'(1'H)-one; 5'-(trifluoromethyl)spiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one; and

5',6'-dimethylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one.

In one embodiment, the present invention concerns also the synthesis

intermediates of formula VI or stereoisomeric forms thereof,




wherein

R¹ is hydrogen,


R² is hydrogen or C1-20-alkyl,

R³ is hydrogen, C1-20-alkyl, C4-8-cycloalkyl, C5-8-cycloalkenyl, aryl, aromatic or non aromatic heterocycle, C1-20-alkoxy, or a group of formula -W-R⁸,

5 R^{3a} is hydrogen, C1-20-alkyl or a group of formula:



or NR³R^{3a} is a group of formula:



10 R⁴ is hydrogen,

R⁵ is hydrogen; halogen; azido; cyano; -S-C1-4-alkyl; -SO-C1-4-alkyl;

-SO₂-C1-4-alkyl; -SONH₂; or C1-20-alkyl unsubstituted or substituted by halogen,

R⁶ is hydrogen, C1-20-alkyl or halogen,

R⁷ is hydrogen, C2-20-alkyl or halogen,

15 W is C1-12-alkylene, -NH- or -NHC(=O)-,

X is O, S or NH,

Y is O, S, -CR¹²R¹³-, -NR¹⁴- or -C(=O)-,

R⁸ is aryl or heterocycle,

R⁹, R¹⁰, R^{10a} and R¹¹ are independently selected from hydrogen, C1-4-alkyl,

20 halogen, hydroxy or methoxycarbonyl,

or R¹⁰ and R^{10a} together form a C3-6-alkylene,

R¹² is hydrogen, C1-4-alkyl, halogen or hydroxy,

R¹³ is hydrogen,

or CR¹²R¹³ is dioxolanyl,

25 R¹⁴ is aryl, heterocycle or a group of formula -V-R¹⁵,

V is C1-12-alkylene,


R¹⁵ is aryl or heterocycle,

m is 1 to 4,

n is 0 or 1,

30 and at least one of R⁵, R⁶ or R⁷ is different from hydrogen when R² is hydrogen, R³ is H or 2,6-diisopropylphenyl, and R^{3a} is H.

In another embodiment, the present invention concerns the synthesis intermediates of formula VI or stereoisomeric forms thereof,




wherein

R¹ is hydrogen,


R² is hydrogen or C1-20-alkyl,

5 R³ is hydrogen, C1-20-alkyl, C4-8-cycloalkyl, C5-8-cycloalkenyl, aryl, aromatic or non aromatic heterocycle, C1-20-alkoxy, or a group of formula -W-R⁸,

R^{3a} is hydrogen, C1-20-alkyl or a group of formula:



or NR³R^{3a} is a group of formula:



10 R⁴ is hydrogen,

R⁵ is hydrogen; halogen; or C1-20-alkyl unsubstituted or substituted by halogen,

R⁶ is hydrogen, C1-20-alkyl or halogen,

R⁷ is hydrogen, C2-20-alkyl or halogen,

15 W is C1-12-alkylene, -NH- or -NHC(=O)-,

X is O, S or NH,

Y is O, S, -CR¹²R¹³-, -NR¹⁴- or -C(=O)-,

R⁸ is aryl or heterocycle,

R⁹, R¹⁰, R^{10a} and R¹¹ are independently selected from hydrogen, C1-4-alkyl,

20 halogen, hydroxy or methoxycarbonyl,

or R¹⁰ and R^{10a} together form a C3-6-alkylene,

R¹² is hydrogen, C1-4-alkyl, halogen or hydroxy,

R¹³ is hydrogen,

or CR¹²R¹³ is dioxolanyl,

25 R¹⁴ is aryl, heterocycle or a group of formula -V-R¹⁵,

V is C1-12-alkylene,

R¹⁵ is aryl or heterocycle,


m is 1 to 4,

n is 0 or 1,

and at least one of R⁵, R⁶ or R⁷ is different from hydrogen when R² is hydrogen, R³ is H or 2,6-diisopropylphenyl, and R^{3a} is H.

Preferably, the synthesis intermediates of formula VI are selected from the group consisting of : 2-(5-chloro-1H-indol-1-yl)propanamide;
2-(7-chloro-1H-indol-1-yl)acetamide; 2-(6-chloro-1H-indol-1-yl)acetamide,
2-(5-chloro-1H-indol-1-yl)butanamide; 2-(5-methyl-1H-indol-1-yl)propanamide; 2-(5-bromo-1H-indol-1-yl)propanamide; 2-(7-fluoro-1H-indol-1-yl)acetamide; 2-(5-bromo-1H-indol-1-yl)acetamide; 2-(5-fluoro-1H-indol-1-yl)acetamide; and 2-(5-chloro-1H-indol-1-yl)acetamide.


In one embodiment, the present invention concerns also the synthesis intermediates of formula IX or stereoisomeric forms thereof,



wherein

- 15 R¹ is hydrogen,
R² is hydrogen or C1-20-alkyl,
R⁴ is hydrogen,
R⁵ is hydrogen; nitro; halogen; azido; cyano; -S-C1-4-alkyl; -SO-C1-4-alkyl;
-SO₂-C1-4-alkyl; -SONH₂; C1-20-alkyl unsubstituted or substituted by halogen; or
20 C1-20-alkoxy unsubstituted or substituted by halogen,
R⁶ is hydrogen, C1-20-alkyl or halogen,
R⁷ is hydrogen, C1-20-alkyl or halogen,
and at least one of R⁵, R⁶ or R⁷ is different from hydrogen when R² is hydrogen, R³ is H or 2,6-diisopropylphenyl, and R^{3a} is H.

25 In another embodiment, the present invention concerns the synthesis intermediates of formula IX or stereoisomeric forms thereof,



wherein

R¹ is hydrogen,

R² is hydrogen or C1-20-alkyl,

R⁴ is hydrogen,

R⁵ is hydrogen; nitro; halogen; C1-20-alkyl unsubstituted or substituted by halogen; or C1-20-alkoxy unsubstituted or substituted by halogen,

5 R⁶ is hydrogen, C1-20-alkyl or halogen,

R⁷ is hydrogen, C1-20-alkyl or halogen,

and at least one of R⁵, R⁶ or R⁷ is different from hydrogen when R² is hydrogen, R³ is H or 2,6-diisopropylphenyl, and R^{3a} is H.

10 Preferably, the synthesis intermediate of formula IX is (5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetic acid.

The present invention also concerns the synthesis intermediates 2-(7-fluoro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide and ethyl (5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetate.

15 It has now been found that compounds of formula I and their pharmaceutically acceptable salts are useful in a variety of pharmaceutical indications.

For example, the compounds according to the invention are useful for the treatment of epilepsy, epileptogenesis, seizure disorders and convulsions.

These compounds may also be used for the treatment of Parkinson's disease.

20 These compounds may also be used for the treatment of dyskinesia induced by dopamine replacement therapy, tardive dyskinesia induced by administration of neuroleptic drugs or Huntington Chorea.

25 In addition, the compounds according to the invention may also be used for treating other neurological disorders including bipolar disorders, mania, depression, anxiety, attention deficit hyperactivity disorder (ADHD), migraine, trigeminal and other neuralgia, chronic pain, neuropathic pain, cerebral ischemia, cardiac arrhythmia, myotonia, cocaine abuse, stroke, myoclonus, tremor, essential tremor, simple or complex tics, Tourette syndrome, restless leg syndrome and other movement disorders, neonatal cerebral haemorrhage, amyotrophic lateral sclerosis, spasticity and degenerative diseases, bronchial asthma, asthmatic status and allergic bronchitis, 30 asthmatic syndrome, bronchial hyperreactivity and bronchospastic syndromes as well as allergic and vasomotor rhinitis and rhinoconjunctivitis.

Thus, the present invention also concerns a compound having the formula I or a pharmaceutically acceptable salt thereof or stereoisomeric forms thereof as defined above for use as a medicament.

35 In a further aspect, the present invention concerns also the use of a compound of formula I or a pharmaceutically acceptable salt thereof for the manufacture of a

medicament for the treatment of neurological and other disorders such as mentioned above.

In particular, the present invention concerns the use of a compound of formula I or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of epilepsy, Parkinson's disease, dyskinesia, migraine, bipolar disorders, chronic pain, neuropathic pain, or bronchial, asthmatic or allergic conditions.

The methods of the invention comprise administration to a mammal (preferably human) suffering from above mentioned conditions or disorders, of a compound according to the invention in an amount sufficient to alleviate or prevent the disorder or condition.

The compound is conveniently administered in any suitable unit dosage form, including but not limited to one containing 3 to 3000 mg, preferably 25 to 500 mg of active ingredient per unit dosage form.

The term "treatment" as used herein includes curative treatment and prophylactic treatment.

By "curative" is meant efficacy in treating a current symptomatic episode of a disorder or condition.

By "prophylactic" is meant prevention of the occurrence or recurrence of a disorder or condition.

The term "epilepsy" as used herein refers to a chronic neurologic condition characterised by unprovoked, recurrent epileptic seizures. An epileptic seizure is the manifestation of an abnormal and excessive synchronised discharge of a set of cerebral neurons; its clinical manifestations are sudden and transient. The term "epilepsy" as used herein can also refer to a disorder of brain function characterised by the periodic occurrence of seizures. Seizures can be "nonepileptic" when evoked in a normal brain by conditions such as high fever or exposure to toxins or "epileptic" when evoked without evident provocation.

The term "seizure" as used herein refers to a transient alteration of behaviour due to the disordered, synchronous, and rhythmic firing of populations of brain neurones.

The term "Parkinsonian symptoms" relates to a syndrome characterized by slowness of movement (bradykinesia), rigidity and / or tremor. Parkinsonian symptoms are seen in a variety of conditions, most commonly in idiopathic Parkinsonism (i.e. Parkinson's Disease) but also following treatment of schizophrenia, exposure to toxins/drugs and head injury. It is widely appreciated that the primary pathology underlying Parkinson's disease is degeneration, in the brain, of the

dopaminergic projection from the substantia nigra to the striatum. This has led to the widespread use of dopamine-replacing agents (e.g. L-3,4-dihydroxyphenylalanine (L-DOPA) and dopamine agonists) as symptomatic treatments for Parkinson's disease and such treatments have been successful in increasing the quality of life of patients suffering from Parkinson's disease. However, dopamine-replacement treatments do have limitations, especially following long-term treatment. Problems can include a wearing-off of the anti-parkinsonian efficacy of the treatment and the appearance of a range of side-effects which manifest as abnormal involuntary movements, such as dyskinesias.

The term "dyskinesia" is defined as the development in a subject of abnormal involuntary movements. This appears in patients with Huntington's disease, in Parkinson's disease patients exposed to chronic dopamine replacement therapy, and in Schizophrenia patients exposed to chronic treatment with neuroleptics. Dyskinesias, as a whole, are characterised by the development in a subject of abnormal involuntary movements. One way in which dyskinesias may arise is as a side effect of dopamine replacement therapy for parkinsonism or other basal ganglia-related movement disorders.

The term "migraine" as used herein means a disorder characterised by recurrent attacks of headache that vary widely in intensity, frequency, and duration.

The attacks are commonly unilateral and are usually associated with anorexia, nausea, vomiting, phonophobia, and/or photophobia. In some cases they are preceded by, or associated with, neurological and mood disturbances. Migraine headache may last from 4 hours to about 72 hours. The International Headache Society (IHS, 1988) classifies migraine with aura (classical migraine) and migraine without aura (common migraine) as the major types of migraine. Migraine with aura consists of a headache phase preceded by characteristic visual, sensory, speech, or motor symptoms. In the absence of such symptoms, the headache is called migraine without aura.

The term "bipolar disorders" as used herein refers to those disorders classified as Mood Disorders according to the Diagnostic and Statistical Manual of Mental

Disorders, 4th edition (Diagnostic and Statistical Manual of Mental Disorders (DSM-IV TM), American Psychiatry Association, Washington, DC, 1994). Bipolar disorders are generally characterised by spontaneously triggered repeated (i.e. at least two) episodes in which the patient's hyperexcitability, activity and mood are significantly disturbed, this disturbance consisting on some occasions of an elevation of mood and increased energy and activity (mania or hypomania), and in other occasions a lowering of mood and decreased energy and activity (depression). Bipolar disorders are separated into four main categories in the DSM-IV (bipolar I disorder, bipolar II disorder,

cyclothymia, and bipolar disorders not otherwise specified).

The term "manic episode", as used herein refers to a distinct period during which there is an abnormally and persistently elevated, expansive, or irritable mood with signs of pressured speech and psychomotor agitation.

5 The term "hypomania", as used herein refers to a less extreme manic episode, with lower grade of severity.

The term "major depressive episode", as used herein refers to a period of at least 2 weeks during which there is either depressed mood or the loss of interest or pleasure in nearly all activities with signs of impaired concentration and psychomotor
10 retardation.

The term "mixed episode", as used herein refers to a period of time (lasting at least 1 week) in which the criteria are met both for a manic episode and for a major depressive episode nearly every day.

The term "chronic pain" as used herein refers to the condition gradually being
15 recognised as a disease process distinct from acute pain. Conventionally defined as pain that persists beyond the normal time of healing, pain can also be considered chronic at the point when the individual realises that the pain is going to be a persistent part of their lives for the foreseeable future. It is likely that a majority of chronic pain syndromes involves a neuropathic component, which is usually harder to
20 treat than acute somatic pain.

The term "neuropathic pain" as used herein refers to pain due to a dysfunctional nervous system, sometimes occurring following injury to the central nervous system (central pain), but more often caused by damage to peripheral nerves (painful peripheral neuropathy). Neuropathic pain is most likely caused by neural
25 hyperexcitation in partially damaged nerves. Several types of painful peripheral neuropathy, which may share some underlying pathogenic mechanisms, have been distinguished, such as: (1) posttraumatic painful peripheral neuropathy; (2) phantom limb pain; (3) facial (trigeminal) pains; (4) postherpetic neuralgia; (5) painful diabetic neuropathy; (6) neuropathies due to cancer tumors; (7) neuropathies induced by
30 treatment with anti-neoplastic agents; and (8) nerve damage associated with demyelinating disease, such as multiple sclerosis. In neuropathic pain, painful reactions appear in response to normally neutral stimuli (allodynia) or as exaggerated reactions to painful stimuli (hyperalgesia). Spontaneous pain, not provoked by external stimuli, also occurs in neuropathic pain, and is the most difficult form of pain
35 to measure and treat.

The term "tics" refers to common and often disabling neurological disorders. They are frequently associated with behaviour difficulties, including obsessive-

compulsive disorder, attention deficit hyperactivity disorder and impulse control. Tics are involuntary, sudden, rapid, repetitive, nonrhythmic stereotype movements or vocalizations. Tics are manifested in a variety of forms, with different durations and degrees of complexity. Simple motor tics are brief rapid movements that often involve only one muscle group. Complex motor tics are abrupt movements that involve either a cluster of simple movements or a more coordinated sequence of movements. Simple vocal tics include sounds such as grunting, barking, yelping, and throat clearing. Complex vocal tics include syllables, phrases, repeating other people's words and repeating one's own words.

The term "tremor" refers to an involuntary, rhythmical, oscillatory movement of a body part. Tremor can be phenomenologically defined as tremor at rest or associated with an action. Such an action can be postural (maintenance of a limb position), kinetic (movement-related), or intentional (at the end of a purposeful movement). Etiologically, tremor most often occurs in Parkinson's disease (Parkinsonian rest tremor) and in essential tremor (postural and kinetic tremor), which consists of hereditary and age-related forms. Tremor may also occur in dystonia and in multiple sclerosis. Other tremors, which can arise from various etiologies, are cerebellar (intentional tremor) and Holmes' midbrain tremor (postural tremor). Tremor can also be an exaggerated form of normal physiological tremor. Apart from the behavioral context in which tremor occurs, tremor frequency is an important criterion to distinguish between various forms of tremor. Essential tremor has the highest incidence of all tremors. As it is age-related, it can be expected to increase in aging populations. Animal model and clinical data indicate that essential tremor may be primarily based on a brainstem (inferior olivary nucleus) - cerebellar dysfunction, whereas Parkinsonian tremor probably originates from abnormal activity within the basal ganglia. Excessive synchronization and/or hyperexcitation in neuronal circuits may underlie tremor activity.

The activity of the compounds of formula I, or their pharmaceutically acceptable salts, as anticonvulsants can be determined in the audiogenic seizures model. The objective of this test is to evaluate the anticonvulsant potential of a compound by means of audiogenic seizures induced in sound-susceptible mice, a genetic animal model with reflex seizures. In this model of primary generalised epilepsy, seizures are evoked without electrical or chemical stimulation and the seizure types are, at least in part, similar in their clinical phenomenology to seizures occurring in man (Löscher W. & Schmidt D., *Epilepsy Res.* (1998), 2, p. 145-181; Buchhalter J.R., *Epilepsia* (1993), 34, S31-S41). Results obtained with compounds of formula I are indicative of a strong pharmacological effect.

Another assay indicative of potential anticonvulsant activity is binding to levetiracetam binding site (LBS) as hereinafter described.

Activity in any of the above-mentioned indications can of course be determined by carrying out suitable clinical trials in a manner known to a person skilled in the relevant art for the particular indication and/or in the design of clinical trials in general.

For treating diseases, compounds of formula I or their pharmaceutically acceptable salts may be employed at an effective daily dosage and administered in the form of a pharmaceutical composition.

Therefore, another embodiment of the present invention concerns a pharmaceutical composition comprising an effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof in combination with a pharmaceutically acceptable diluent or carrier.

To prepare a pharmaceutical composition according to the invention, one or more of the compounds of formula I or a pharmaceutically acceptable salt thereof is intimately admixed with a pharmaceutical diluent or carrier according to conventional pharmaceutical compounding techniques known to the skilled practitioner.

Suitable diluents and carriers may take a wide variety of forms depending on the desired route of administration, e.g., oral, rectal, parenteral or intranasal.

Pharmaceutical compositions comprising compounds according to the invention can, for example, be administered orally or parenterally, i.e., intravenously, intramuscularly or subcutaneously, intrathecally, by inhalation or intranasally.

Pharmaceutical compositions suitable for oral administration can be solids or liquids and can, for example, be in the form of tablets, pills, dragees, gelatin capsules, solutions, syrups, chewing-gums and the like.

To this end the active ingredient may be mixed with an inert diluent or a non-toxic pharmaceutically acceptable carrier such as starch or lactose. Optionally, these pharmaceutical compositions can also contain a binder such as microcrystalline cellulose, gum tragacanth or gelatine, a disintegrant such as alginic acid, a lubricant such as magnesium stearate, a glidant such as colloidal silicon dioxide, a sweetener such as sucrose or saccharin, or colouring agents or a flavouring agent such as peppermint or methyl salicylate.

The invention also contemplates compositions which can release the active substance in a controlled manner. Pharmaceutical compositions which can be used for parenteral administration are in conventional form such as aqueous or oily solutions or suspensions generally contained in ampoules, disposable syringes, glass or plastics vials or infusion containers.

In addition to the active ingredient, these solutions or suspensions can optionally also contain a sterile diluent such as water for injection, a physiological saline solution, oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents, antibacterial agents such as benzyl alcohol, antioxidants such as ascorbic acid or sodium bisulphite, chelating agents such as ethylene diamine-tetra-acetic acid, buffers such as acetates, citrates or phosphates and agents for adjusting the osmolality, such as sodium chloride or dextrose.

These pharmaceutical forms are prepared using methods which are routinely used by pharmacists.

The amount of active ingredient in the pharmaceutical compositions can fall within a wide range of concentrations and depends on a variety of factors such as the patient's sex, age, weight and medical condition, as well as on the method of administration. Thus the quantity of compound of formula I in compositions for oral administration is at least 0.5 % by weight and can be up to 80 % by weight with respect to the total weight of the composition.

In accordance with the invention it has also been found that the compounds of formula I or the pharmaceutically acceptable salts thereof can be administered alone or in combination with other pharmaceutically active ingredients. Non-limiting examples of such additional compounds which can be cited for use in combination with the compounds according to the invention are antivirals, antispastics (e.g. baclofen), antiemetics, antimanic mood stabilizing agents, analgesics (e.g. aspirin, ibuprofen, paracetamol), narcotic analgesics, topical anesthetics, opioid analgesics, lithium salts, antidepressants (e.g. mianserin, fluoxetine, trazodone), tricyclic antidepressants (e.g. imipramine, desipramine), anticonvulsants (e.g. valproic acid, carbamazepine, phenytoin), antipsychotics (e.g. risperidone, haloperidol), neuroleptics, benzodiazepines (e.g. diazepam, clonazepam), phenothiazines (e.g. chlorpromazine), calcium channel blockers, amphetamine, clonidine, lidocaine, mexiletine, capsaicin, caffeine, quetiapine, serotonin antagonists, β -blockers, antiarrhythmics, triptans, ergot derivatives and amantadine.

Of particular interest in accordance with the present invention are combinations of at least one compound of formula I or a pharmaceutically acceptable salt thereof and at least one compound inducing neural inhibition mediated by GABA_A receptors. The compounds of formula I exhibit a potentiating effect on the compounds inducing neural inhibition mediated by GABA_A receptors enabling, in many cases, effective treatment of conditions and disorders under reduced risk of adverse effects.

Examples of compounds inducing neural inhibition mediated by GABA_A receptors include the following: benzodiazepines, barbiturates, steroids, and

anticonvulsants such as valproate, vigabatrine, tiagabine or pharmaceutical acceptable salts thereof.

5 Benzodiazepines include the 1,4-benzodiazepines, such as diazepam and clonazepam, and the 1,5-benzodiazepines, such as clobazam. Preferred compound is clonazepam.

Barbiturates include phenobarbital and pentobarbital. Preferred compound is phenobarbital.

Steroids include adrenocorticotrophic hormones such as tetracosactide acetate, etc.

10 Anticonvulsants include hydantoin (phenytoin, ethotoin, etc), oxazolindines (trimethadione, etc.), succinimides (ethosuximide, etc.), phenacemides (phenacemide, acetylpheneturide, etc.), sulfonamides (sulthiame, acetoazolamide, etc.), aminobutyric acids (e.g. gamma-amino-beta-hydroxybutyric acid, etc.), sodium valproate and derivatives, carbamazepine and so on.

15 Preferred compounds include valproic acid, valpromide, valproate pivoxil, sodium valproate, semi-sodium valproate, divalproex, clonazepam, phenobarbital, vigabatrine, tiagabine and amantadine.

Of particular interest in accordance with the present invention are combinations of at least one compound of formula I or a pharmaceutically acceptable salt thereof and at least one compound modulating neurotransmission mediated by glutamate receptors. The compounds of formula I exhibit a potentiating effect on the compounds modulating neurotransmission mediated by glutamate receptors enabling, in many cases, effective treatment of conditions and disorders under reduced risk of adverse effects.

25 Examples of compounds modulating neurotransmission mediated by glutamate receptors include the following: NBQX and MK-801 or pharmaceutical acceptable salts thereof.

For the preferred oral compositions, the daily dosage is in the range 3 to 3000 milligrams (mg) of compounds of formula I.

30 In compositions for parenteral administration, the quantity of compound of formula I present is at least 0.5 % by weight and can be up to 33 % by weight with respect to the total weight of the composition. For the preferred parenteral compositions, the dosage unit is in the range 3 mg to 3000 mg of compounds of formula I.

35 The daily dose can fall within a wide range of dosage units of compound of formula I and is generally in the range 3 to 3000 mg. However, it should be

understood that the specific doses can be adapted to particular cases depending on the individual requirements, at the physician's discretion.

The LBS binding compounds provided by this invention and labelled derivatives thereof may be useful as standards and reagents in determining the ability of tested compounds (e.g., a potential pharmaceutical) to bind to the LBS receptor.

Labelled derivatives of LBS ligands provided by this invention may also be useful as radiotracers for positron emission tomography (PET) imaging or for single photon emission computerized tomography (SPECT).

The following examples are provided for illustrative purposes.

Unless specified otherwise in the examples, characterization of the compounds is performed according to the following methods:

NMR spectra are recorded on a BRUKER AC 250 Fourier Transform NMR Spectrometer fitted with an Aspect 3000 computer and a 5mm $^1\text{H}/^{13}\text{C}$ dual probehead or BRUKER DRX 400 FT NMR fitted with a SG Indigo² computer and a 5 mm inverse geometry $^1\text{H}/^{13}\text{C}/^{15}\text{N}$ triple probehead. The compound is studied in DMSO- d_6 (or CDCl_3) solution at a probe temperature of 313 K or 300 K and at a concentration of 20 mg/ml. The instrument is locked on the deuterium signal of DMSO- d_6 (or CDCl_3). Chemical shifts are given in ppm downfield from TMS taken as internal standard.

HPLC analyses are performed using one of the following systems:

- an Agilent 1100 series HPLC system mounted with an INERTSIL ODS 3 C18, DP 5 μm , 250 X 4.6 mm column. The gradient runs from 100 % solvent A (acetonitrile, water, H_3PO_4 (5/95/0.001, v/v/v)) to 100 % solvent B (acetonitrile, water, H_3PO_4 (95/5/0.001, v/v/v)) in 6 min with a hold at 100 % B of 4 min. The flow rate is set at 2.5 ml/min. The chromatography is carried out at 35°C.

- a HP 1090 series HPLC system mounted with a HPLC Waters Symetry C18, 250 X 4.6 mm column. The gradient runs from 100 % solvent A (MeOH, water, H_3PO_4 (15/85/0.001M, v/v/M)) to 100 % solvent B (MeOH, water, H_3PO_4 (85/15/0.001 M, v/v/M)) in 10 min with a hold at 100 % B of 10 min. The flow rate is set at 1 ml/min. The chromatography is carried out at 40 °C.

Mass spectrometric measurements in LC/MS mode are performed as follows:

HPLC conditions

Analyses are performed using a WATERS Alliance HPLC system mounted with an INERTSIL ODS 3, DP 5 μm , 250 X 4.6 mm column.

The gradient runs from 100 % solvent A (acetonitrile, water, TFA (10/90/0.1, v/v/v)) to 100 % solvent B (acetonitrile, water, TFA (90/10/0.1, v/v/v)) in 7 min with a

hold at 100 % B of 4 min. The flow rate is set at 2.5 ml/min and a split of 1/25 is used just before API source.

MS conditions

Samples are dissolved in acetonitrile/water, 70/30, v/v at the concentration of
5 about 250 µgr/ml. API spectra (+ or -) are performed using a FINNIGAN (San Jose, CA, USA) LCQ ion trap mass spectrometer. APCI source operates at 450 °C and the capillary heater at 160 °C. ESI source operates at 3.5 kV and the capillary heater at 210 °C.

Mass spectrometric measurements in DIP/EI mode are performed as follows:
10 samples are vaporized by heating the probe from 50 °C to 250 °C in 5 min. EI (Electron Impact) spectra are recorded using a FINNIGAN (San Jose, CA, USA) TSQ 700 tandem quadrupole mass spectrometer. The source temperature is set at 150 °C.

Mass spectrometric measurements on a TSQ 700 tandem quadrupole mass spectrometer (Finnigan MAT, San Jose, CA, USA) in GC/MS mode are performed with
15 a gas chromatograph model 3400 (Varian, Walnut Creek, CA, USA) fitted with a split/splitless injector and a DB-5MS fused-silica column (15 m x 0.25 mm I.D., 1 µm) from J&W Scientific (Folsom, CA, USA). Helium (purity 99.999 %) is used as carrier gas. The injector (CTC A200S autosampler) and the transfer line operate at 290 and 250 °C, respectively. Sample (1 µl) is injected in splitless mode and the oven
20 temperature is programmed as follows: 50 °C for 5 min., increasing to 280 °C (23°C/min) and holding for 10 min. The TSQ 700 spectrometer operates in electron impact (EI) or chemical ionization (CI/CH₄) mode (mass range 33 - 800, scan time 1.00 sec). The source temperature is set at 150 °C.

Specific rotation is recorded on a Perkin-Elmer 341 polarimeter. The angle of
25 rotation is recorded at 25 °C on 1 % solutions in MeOH. For some molecules, the solvent is CH₂Cl₂ or DMSO, due to solubility problems.

Melting points are determined on a Büchi 535 or 545 Tottoli-type fusionometre, and are not corrected, or by the onset temperature on a Perkin Elmer DSC 7.

30 *Preparative chromatographic separations* are performed on silicagel 60 Merck, particle size 15-40 µm, reference 1.15111.9025, using Novasep axial compression columns (80 mm i.d.), flow rates between 70 and 150 ml/min. Amount of silicagel and solvent mixtures as described in individual procedures.


Preparative Chiral Chromatographic separations are performed on a DAICEL
35 Chiralpak AD 20 µm, 100*500 mm column using an in-house build instrument with various mixtures of lower alcohols and C5 to C8 linear, branched or cyclic alkanes at ± 350 ml/min. Solvent mixtures as described in individual procedures.

The following abbreviations are used in the examples:

AcOEt	Ethyl acetate
CH ₃ CN	Acetonitrile
DMF	N,N-Dimethylformamide
5 NBS	N-bromosuccinimide
NCS	N-chlorosuccinimide
NIS	N-iodosuccinimide
TFA	Trifluoroacetic acid
10 THF	Tetrahydrofuran

In the tables, the stereochemical information is contained in the two columns headed "configuration". The second column indicates whether a compound has no stereogenic center (achiral), is a pure enantiomer (pure), a racemate (rac) or is a mixture of two stereoisomers, possibly in unequal proportions (MIXT). The first column contains the stereochemical assignment for the recognised center, following the IUPAC numbering used in the "IUPAC name" column. A number alone indicates the existence of both configurations at that center. A number followed by 'R' or 'S' indicates the known absolute configuration at that center. A number followed by '§' indicates the existence of only one but unknown absolute configuration at that center. The letter (A, B) in front is a way of distinguishing the various enantiomers of the same structure.

Example 1: Synthesis of 2-(5-iodo-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide **2**.



2-(2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide **1** is synthesized according to the method described by Valenta et al. (Valenta, V.; Holubeck, J.; Svatek, E.; Valchar, M.; Krejci, I.; Protiva, M.; Collect. Czech. Chem. Commun. (1990), 55, 2756-2764).

Oxindole **1** (1 g, 5.25 mmol) was dissolved in CH₃CN (20 ml). After addition of the NIS (1.3 g, 5.78 mmol), the TFA (217 μ l, 1.57 mmol) was added and the reaction was allowed at room temperature for 16 h. After evaporation of the solvent, the mixture was triturated in a 10 % aqueous solution of Na₂S₂O₃. The beige solid formed was filtered, washed with water and with ether. After crystallization from 90 %


aqueous EtOH, and re-crystallization from acetonitrile, 2-(5-iodo-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide **2** was obtained as a white solid.

Yield: 166 mg (10 %).

MS (GC-MS, M^+): 316.

5

Example 2: Synthesis of 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide **3**.



10

Oxindole **1** (1.77 g, 9.3 mmol) was dissolved in 90 % H_2SO_4 (6 ml) at room temperature, and NCS (1.24 g, 9.3 mmol) was slowly added with stirring. After 2 hours, the mixture was poured into cold water. The precipitate was collected, washed several times with water and then with Et_2O . After crystallization from EtOH, 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide **3** was obtained as a white solid.


Yield: 479 mg (23 %).

15

MS (LC-MS, MH^+): 225/227.

MP: 226 °C.

Example 3: Synthesis of 2-(5,7-dibromo-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide **4**.



20

2-(5,7-dibromo-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide **4** was obtained as described in example 1 by using 2 equivalents of NBS. The crude material was purified by silica gel chromatography.


25

Yield: 129 mg (10 %).

MS (LC-MS, MH^+): 259/261.

Example 4: Synthesis of 2-(5-nitro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide **5**.

35




To a stirred solution of 2-(2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide **1** (400 mg, 2.1 mmol) in TFA (20 ml) was added fuming nitric acid (170 μ l, 2.7 mmol) over 10 minutes. Following addition, the ice bath was removed and the mixture was stirred at room temperature for 5 minutes, then poured carefully into ice water. The precipitate was collected, washed with water until pH 7 and dried to give a crude solid. Crystallization in a mixture acetonitrile/MeOH afforded the 2-(5-nitro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide **5** as a green-gray solid.

Yield: 150 mg (30 %).

MS (DIP, M^+): 235.

Example 5: Synthesis of 2-(5-methyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide **24**.



5.1 Synthesis of 5'-methylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one **6**.

5-methyl-1H-indole-2,3-dione (6 g, 37 mmol) was suspended in 100 ml of AcOH. The heterogeneous mixture was heated at 60 °C. After complete solubilisation, 1,2-ethanedithiol (3.15 ml, 37 mmol) was added and then neat $\text{BF}_3 \cdot \text{OEt}_2$ (9.5 ml, 75 mmol) was added dropwise. The reaction was stirred for 25 minutes, in which time the reaction mixture became homogenous. After 20 minutes at room temperature, the reaction was quenched by addition of water, the solid washed several times with large amounts of water and air dried affording 5'-methylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one **6** as a brown solid.

Yield: 8.65 g (98 %).

MS (DIP, M^+): 237.

Compounds listed in table 1 can be synthesised according to the same method.

Table 1

No	IUPAC NAME
6	5'-methylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one
7	5'-[(trifluoromethyl)oxy]spiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one
8	5'-(1-methylethyl)spiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one
9	5'-ethylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one
10	5'-fluorospiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one
11	5',7'-dimethylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one
12	5'-propylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one
13	5'-(trifluoromethyl)spiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one
14	5',6'-dimethylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one

5.2 Synthesis of 2-[5'-methyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl]acetamide **15**.

Compound **6** (8 g, 33.7 mmol) was dissolved in dry DMF (80 ml) under a nitrogen atmosphere. The solution was cooled at 0 °C and NaH (1.62 g, 37.13 mmol, 60 % dispersion) was carefully added portionwise. When the nitrogen evolution ceased, bromoacetamide (5.6 g, 37.13 mmol) was added. After 30 minutes, the mixture was poured into cold water and the solid filtered off, washed with water and hexane. The crude material was directly crystallized in acetonitrile affording 2-[5'-methyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl]acetamide **15** as a white solid.

Yield: 4.86 g (49 %).

MS (LC-MS, MH⁺): 295.

Compounds listed in table 2 can be synthesised according to the same method.

Table 2:

No	IUPAC NAME
15	2-(5'-methyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide
16	2-[2'-oxo-5'-[(trifluoromethyl)oxy]spiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl]acetamide
17	2-[5'-(1-methylethyl)-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl]acetamide
18	2-(5'-ethyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide
19	2-(5'-fluoro-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide
20	2-(5',7'-dimethyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide
21	2-(2'-oxo-5'-propylspiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide
22	2-[2'-oxo-5'-(trifluoromethyl)spiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl]acetamide
23	2-(5',6'-dimethyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide


5.3 Synthesis of 2-(5-methyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide **24**.

The Raney Nickel was prepared as an aqueous slurry after removing four fifth
of water of the commercial solution. Aqueous Raney nickel (10 ml) was added to a
solution of compound **15** (4.06 g, 13.8 mmol) in 40 ml of distilled THF and the mixture
was further vigorously stirred at room temperature. When no starting material was
detected by thin layer chromatography, the mixture was diluted with THF and filtered
through a Celite pad. After removal of the solvent, the crude material was purified by
silica gel chromatography (CH₂Cl₂/MeOH 95/5 then 90/10), and the solvent was
evaporated to yield the 2-(5-methyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide **24** as a
white solid.

Yield: 697 mg (21 %).

MS (LC-MS, MH⁺): 205.

Example 6: Synthesis of 2-(5-chloro-2-oxo-2,3-dihydro-1*H*-indol-1-yl)propanamide **32**, **33** and **34**.



5

6.1 Synthesis of 2-(5-chloro-1*H*-indol-1-yl)propanamide **25**.

A dispersion of 60 % NaH (6.85 g, 0.17 mol) was added to an ice-cooled solution of 5-chloroindole (20 g, 0.13 mol) in 250 ml of dry DMF. The stirring was continued for 20 minutes at room temperature, and the mixture was cooled again with an ice bath. After portionwise addition of solid 2-bromopropanamide (24.1 g, 0.15 mol), the reaction mixture was stirred for 1h30 at room temperature, then poured into cold water and extracted 3 times with AcOEt. The combined organic phases were dried over Na₂SO₄, filtered and concentrated in vacuo. The crude material was purified by silica gel chromatography (AcOEt/hexane 50/50) to give pure 2-(5-chloro-1*H*-indol-1-yl)propanamide **25** as a white solid.

Yield: 13.45 g (46 %).

MS (LC-MS, MH⁺): 223/225.

Compounds listed in table 3 can be synthesised according to the same method.

Table 3:

No	Configuration	IUPAC NAME
25	2 rac	2-(5-chloro-1H-indol-1-yl)propanamide
26	achiral	2-(7-chloro-1H-indol-1-yl)acetamide
27	achiral	2-(6-chloro-1H-indol-1-yl)acetamide
28	2 rac	2-(5-chloro-1H-indol-1-yl)butanamide
29	2 rac	2-(5-methyl-1H-indol-1-yl)propanamide
30	2 rac	2-(5-bromo-1H-indol-1-yl)propanamide
55	achiral	2-(7-fluoro-1H-indol-1-yl)acetamide
112	achiral	2-(5-bromo-1H-indol-1-yl)acetamide
113	achiral	2-(5-fluoro-1H-indol-1-yl)acetamide
114	achiral	2-(5-chloro-1H-indol-1-yl)acetamide

6.2 Synthesis of 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide **32**.

5 Pyridinium bromide perbromide (38.8 g, 117 mmol) was added in portions over a period of 30 minutes to a stirred solution of 2-(5-chloro-1H-indol-1-yl)propanamide 25 (13 g, 58.4 mmol) in *tert*-butanol (100 ml) at room temperature. The reaction mixture was stirred for 30 minutes, then poured into water and diluted with AcOEt. After removal of the organic layer, the aqueous phase was extracted twice with AcOEt.

10 Combined organic phases were dried over Na₂SO₄ and concentrated. 2-(3,3-dibromo-5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide **31** was obtained as a crude oil and was directly used in the next step, without further purification.

Zinc dust (23.71 g, 0.58 mol) was added to a stirred solution of compound **31** (theoretical: 58.4 mmol) in AcOH (110 ml) at 0 °C. After 1 hour, the reaction mixture

15 was filtered through a Celite pad. The filtrate was diluted with AcOEt and cold water. The pH was adjusted to 7 and the layers were separated. The aqueous phase was extracted again with AcOEt. Organic layers were dried over Na₂SO₄ and concentrated. The beige solid was cristallized in AcOEt and afforded 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide **32** as a white solid.

20 Yield: 2.5 g (18 %).

MS (LC-MS, MH⁺): 239/241.

Compound **32** (2.5 g, 10.5 mmol) was resolved into its enantiomers by chiral chromatography (DAICEL, Chiralcel OD phase, eluent : 50/50 ethanol/hexane) to afford enantiomers **33** (first eluted) and **34** (second eluted) as white solids.

25 Compound **33**:

Yield: 977 mg (39 %).

MS (LC-MS, MH^+): 239/241.

MP: 171-172 °C.


Compound **34**:

Yield: 941 mg (37 %).

5 MS (LC-MS, MH^+): 239/241.

MP: 171-172 °C.

Example 7: Synthesis of N-(4-tert-Butyl-phenyl)-2-(5-chloro-2-oxo-2,3-dihydro-indol-1-yl)-acetamide **85**.



10

7.1 Synthesis of ethyl (5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetate **58**.

The ester **58** was obtained from 5-chloroindole and ethyl bromoacetate (instead of bromo-acetamide) using the methodology described in example 6.

MS (GC-MS, M^+): 253/255.

15

7.2 Synthesis of (5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetic acid **59**.

In a two neck flask, ethyl (5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetate **58** (0.1 g, 0.39 mmol) was stirred overnight at room temperature in aqueous HCl (6 M, 2 ml) and warmed at 80 °C until disappearance of the starting material by TLC. The reaction mixture was cooled down to room temperature, filtered and the filtrate was washed by cold water to afford (5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetic acid **59** (74 mg).

20

LC/MS : 225/227 (MH^+).

25 7.3 Synthesis of N-(4-tert-Butyl-phenyl)-2-(5-chloro-2-oxo-2,3-dihydro-indol-1-yl)-acetamide **85**.

In a 1 ml polypropylene vial, 300 μ l of a 0.15 M stock solution of (5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetic acid **59** in a 1/1 CH_2Cl_2 /DMF mixture were added onto N-dicyclohexyl-N'-methyl polystyrene (from Novabiochem, loading:

1.9 mmol/g, 35 mg) and N,N-diisopropyl-methyl-polystyrene (from Argonaut, loading: 3.49 mmol/g, 25.4 mg) followed by 4-tertbutyl-aniline (0.044 mmol, 6.6 mg). The reaction mixture was stirred 40 h under vortex and quenched with DMF (400 μ l). The suspension was allowed to sediment and the liquid was concentrated under vacuo to afford N-(4-tert-butyl-phenyl)-2-(5-chloro-2-oxo-2,3-dihydro-indol-1-yl)-acetamide **35** (10 mg).

MS (LC-MS, MH^+): 357/359.

Compounds described in table 4 may be prepared according to one of the previous methods.

The synthesis intermediate 2-(7-fluoro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide may also be prepared according to one of the previous methods.

Table 4: Compounds of formula I.

No	Configuration	IUPAC NAME	MS (LC-MS, MH ⁺)	α_D (MeOH, 25 °C, 1 %)
2	achiral	2-(5-iodo-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	316 (GC-MS, M ⁺)	
3	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	225/227	
4	achiral	2-(5,7-dibromo-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	347/349/351	
5	achiral	2-(5-nitro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	235 (DIP, M ⁺)	
24	achiral	2-(5-methyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	205	
32	rac	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide	239/241	
33	pure	(2R)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide	239/241	+64.55
34	pure	(2S)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide	239/241	-60.82
35	achiral	2-[2-oxo-5-(trifluoromethoxy)-2,3-dihydro-1H-indol-1-yl]acetamide	275	
36	achiral	2-(5-isopropyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	233	
37	achiral	2-(5-ethyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	219	
38	achiral	2-(5-fluoro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	209	
39	achiral	2-(5,7-dimethyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	219	
40	achiral	2-(5-bromo-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	269/271	
41	achiral	2-(2-oxo-5-propyl-2,3-dihydro-1H-indol-1-yl)acetamide	233	
42	achiral	2-[2-oxo-5-(trifluoromethyl)-2,3-dihydro-1H-indol-1-yl]acetamide	259	
43	achiral	2-(5,6-dimethyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	219	
44	achiral	2-(7-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	224/226	

No	Configuration	IUPAC NAME	MS (LC-MS, MH ⁺)	α_D (MeOH, 25 °C, 1 %)
45	achiral	2-(6-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	225/227	
46	rac	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)butanamide	252/254	
47	pure	(+)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)butanamide	252/254	+46.23
48	pure	(-)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)butanamide	252/254	-38.68
49	rac	2-(5-methyl-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide	219	
50	pure	(+)-2-(5-methyl-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide	219	+84.39
51	pure	(-)-2-(5-methyl-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide	219	
52	rac	2-(5-bromo-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide	283/285	
53	pure	(-)-2-(5-bromo-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide	283/285	-40.33
54	pure	(+)-2-(5-bromo-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide	283/285	+44.16
57	achiral	2-(5-chloro-7-fluoro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	243/245	
60	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3-hydroxyphenyl)acetamide	317/319	
61	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3-fluorophenyl)acetamide	319/321	
62	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[3-(2-pyridin-2-ylethyl)phenyl]acetamide	406/408	
63	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[6-(hydroxymethyl)cyclohex-3-en-1-yl]acetamide	335/337	

No	Configuration	IUPAC NAME	MS (LC-MS, MH ⁺)	α_D (MeOH, 25 °C, 1 %)
64	achiral	5-chloro-1-[2-oxo-2-(4-pyridin-2-yl)piperazin-1-yl]ethyl]-1,3-dihydro-2H-indol-2-one	371/373	
65	achiral	5-chloro-1-[2-[4-(3-methylphenyl)piperazin-1-yl]-2-oxoethyl]-1,3-dihydro-2H-indol-2-one	384/386	
66	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(4-hydroxy-3-methoxybenzyl)acetamide	361/363	
67	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(pyridin-4-ylmethyl)-N-(tetrahydrofuran-2-ylmethyl)acetamide	400/402	
68	achiral	5-chloro-1-[2-(3-hydroxypiperidin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-indol-2-one	309/311	
69	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N'-isonicotinoylacetohydrazide	345/347	
70	achiral	5-chloro-1-(2-oxo-2-thiomorpholin-4-ylethyl)-1,3-dihydro-2H-indol-2-one	311/313	
71	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(4H-1,2,4-triazol-3-yl)acetamide	292/294	
72	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[4-(methylsulfonyl)benzyl]acetamide	393/395	
73	achiral	1-[5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl]acetyl]octahydroquinolin-4(1H)-one	361/363	
74	achiral	N'-(4-bromophenyl)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetohydrazide	394/396/398	

No	Configuration	IUPAC NAME	MS (LC-MS, MH ⁺)	α_D (MeOH, 25 °C, 1 %)
75	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(6-methoxypyridin-3-yl)acetamide	332/334	
76	achiral	N-butyl-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	281/283	
77	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3-hydroxypropyl)acetamide	283/285	
78	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[3-(dimethylamino)propyl]acetamide	310/312	
79	achiral	5-chloro-1-(2-oxo-2-[4-(2-phenylethyl)piperazin-1-yl]ethyl)-1,3-dihydro-2H-indol-2-one	398/400	
80	achiral	ethyl [(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetyl]amino]acetate	311/313	
81	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3-ethoxypropyl)acetamide	311/313	
82	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(2-fluoroethyl)acetamide	271/273	
83	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-methoxy-N-methylacetamide	269/271	
84	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3,4-dimethylphenyl)acetamide	329/331	
85	achiral	N-(4-tert-butylphenyl)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	357/359	

No	Configuration	IUPAC NAME	MS (LC-MS, MH ⁺)	α_D (MeOH, 25 °C, 1 %)
86	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3-hydroxy-2,2-dimethylpropyl)acetamide	311/313	
87	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[1-(hydroxymethyl)propyl]acetamide	297/299	
88	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3,3,3-trifluoro-2-hydroxypropyl)acetamide	337/339	
89	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(2-hydroxy-2-phenylethyl)acetamide	345/347	
90	achiral	5-chloro-1-[2-[4-(4-hydroxyphenyl)piperazin-1-yl]-2-oxoethyl]-1,3-dihydro-2H-indol-2-one	386/388	
91	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(pyridin-4-ylmethyl)acetamide	316/318	
92	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[(5-methyl-2-furyl)methyl]acetamide	319/321	
93	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[3-(1H-pyrazol-1-yl)propyl]acetamide	333/335	
94	achiral	methyl 3-[(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetyl]-1,3-thiazolidine-4-carboxylate	355/357	
95	achiral	5-chloro-1-[2-(2,5-dihydro-1H-pyrrrol-1-yl)-2-oxoethyl]-1,3-dihydro-2H-indol-2-one	277/279	
96	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N'-(4,5-dihydro-1H-imidazol-2-yl)acetohydrazide	308/310	

No	Configuration	IUPAC NAME	MS (LC-MS, MH ⁺)	α_D (MeOH, 25 °C, 1 %)
97	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-(3,4-dimethoxyphenyl)ethyl]acetamide	389/391	
98	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-(2-chlorophenyl)ethyl]acetamide	363/365/367	
99	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-(4-methylphenyl)ethyl]acetamide	343/345	
100	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(2-morpholin-4-ylethyl)acetamide	338/340	
101	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-(3,4,5,6-tetrahydro-1-benzazocin-1(2H)-yl)propyl]acetamide	426/428	
102	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-(2-methylpiperidin-1-yl)ethyl]acetamide	350/352	
103	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(2-nitrobenzyl)acetamide	360/362	
104	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3,4-dihydro-1H-isochromen-1-ylmethyl)acetamide	371/373	
105	achiral	N-(2-chloro-6-fluorobenzyl)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide	367/369/371	
106	achiral	N-benzyl-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-methylacetamide	329/311	
107	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-[(trifluoromethyl)thio]benzyl]acetamide	415/417	

No	Configuration	IUPAC NAME	MS (LC-MS, MH ⁺)	α_D (MeOH, 25 °C, 1 %)
108	achiral	5-chloro-1-[2-(1,4-dioxo-8-azaspiro[4.5]dec-8-yl)-2-oxoethyl]-1,3-dihydro-2H-indol-2-one	351/353	
109	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-cycloheptylacetamide	321/323	
110	achiral	5-chloro-1-[2-[4-(2-morpholin-4-ylethyl)piperazin-1-yl]-2-oxoethyl]-1,3-dihydro-2H-indol-2-one	407/409	
111	achiral	2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-pyridin-3-ylacetamide	302/304	

Example 8: LBS Binding Assay.

[LBS stands for Levetiracetam Binding Site cf. M. Noyer et al., Eur. J. Pharmacol. (1995), 286, 137-146.]

5 The inhibition constant (K_i) of a compound is determined in competitive binding experiments by measuring the binding of a single concentration of a radioactive ligand at equilibrium with various concentrations of the unlabeled test substance. The concentration of the test substance inhibiting 50 % of the specific binding of the radioligand is called the IC_{50} . The equilibrium dissociation constant K_d is proportional to the IC_{50} and is calculated using the equation of Cheng and Prusoff
10 (Cheng Y. et al., Biochem. Pharmacol. (1972), 22, 3099-3108).

The concentration range usually encompasses 6 log units with variable steps (0.3 to 0.5 log). Assays are performed in mono- or duplicate, each K_i determination is performed on two different samples of test substance.

15 Cerebral cortex from 200-250 g male Sprague-Dawley rats are homogenised using a Potter S homogeniser (10 strokes at 1,000 rpm; Braun, Germany) in 20 mmol/l Tris-HCl (pH 7.4), 250 mmol/l sucrose (buffer A); all operations are performed at 4 °C. The homogenate is centrifuged at 30,000xg for 15 min. The crude membrane pellet obtained is resuspended in 50 mmol/l Tris-HCl (pH 7.4), (buffer B) and incubated 15 min at 37 °C, centrifuged at 30,000xg for 15 min and washed twice with
20 the same buffer. The final pellet is resuspended in buffer A at a protein concentration ranging from 15 to 25 mg/ml and stored in liquid nitrogen.

Membranes (150-200 μ g of protein / assay) are incubated at 4 °C for 120 min in 0.5 ml of a 50 mmol/l Tris-HCl buffer (pH 7.4) containing 2 mmol/l $MgCl_2$, 10^{-9} to $2 \cdot 10^{-9}$ mol/l of [3H]-2-[4-(3-azidophenyl)-2-oxo-1-pyrrolidiny]butanamide and
25 increasing concentrations of the test substance. The non specific binding (NSB) is defined as the residual binding observed in the presence of a concentration of reference substance (e.g. 10^{-3} mol/l levetiracetam) that binds essentially all the receptors. Membrane-bound and free radioligands are separated by rapid filtration through glass fiber filters (equivalent to Whatman GF/C or GF/B; VEL, Belgium) pre-soaked in 0.1 % polyethyleneimine and 10^{-3} mol/l levetiracetam to reduce non
30 specific binding. Samples and filters are rinsed by at least 6 ml of 50 mmol/l Tris-HCl (pH 7.4) buffer. The entire filtration procedure does not exceed 10 seconds per sample. The radioactivity trapped onto the filters is counted by liquid scintillation in a β -counter (Tri-Carb 1900 or TopCount 9206, Camberra Packard, Belgium, or any other
35 equivalent counter). Data analysis is performed by a computerized non linear curve fitting method using a set of equations describing several binding models assuming populations of independent non-interacting receptors, which obey to the law of mass.

Example 9: Animal model of sound-susceptible mice.

The aim of this test is to evaluate the anticonvulsant potency of a compound in sound-susceptible mice, a genetic animal model with reflex seizures. In this model of primary generalised epilepsy, seizures are evoked without electrical or chemical stimulation and the seizure types are, at least in part, similar in their clinical phenomenology to seizures occurring in man (Löscher W. & Schmidt D., *Epilepsy Res.* (1998), 2, 145-181; Buchhalter J.R., *Epilepsia* (1993), 34, S31-S41).


Male or female genetically sound-sensitive mice (14-28 g; N=10), derived from a DBA strain originally selected by Dr. Lehmann of the Laboratory of Acoustic Physiology (Paris) and bred in the UCB Pharma Sector husbandry unit since 1978, are used. The experimental design consists of several groups, one group receiving the vehicle control and the other groups different doses of the test-compound. The compounds are administered intraperitoneally 60 minutes before the induction of audiogenic seizures. The range of the doses administered has a logarithmic progression, generally between 1.0×10^{-5} mol/kg and 1.0×10^{-3} mol/kg, but lower or higher doses are tested if necessary.

For testing, the animals are placed in small cages, one mouse per cage, in a sound-attenuated chamber. After a period of orientation of 30 seconds, the acoustic stimulus (90 dB, 10-20 kHz) is delivered for 30 seconds via loudspeakers positioned above each cage. During this interval, the mice are observed and the presence of the 3 phases of the seizure activity namely wild running, clonic and tonic convulsions, is recorded. The proportion of mice protected against wild running, clonic and tonic convulsions, respectively, is calculated.

For active compounds, an ED₅₀ value, i.e. the dose producing 50 % protection relative to the control group, together with 95 % confidence limits, was calculated using a Probit Analysis (SAS/STAT® Software, version 6.09, PROBIT procedure) of the proportions of protected mice for each of the 3 phases of the seizure activity.

Claims

1. A compound having the formula I or a pharmaceutically acceptable salt thereof or stereoisomeric forms thereof,




5 wherein

R^1 is hydrogen,


R^2 is hydrogen or C1-20-alkyl,

R^3 is hydrogen, C1-20-alkyl, C4-8-cycloalkyl, C5-8-cycloalkenyl, aryl, aromatic or non aromatic heterocycle, C1-20-alkoxy, or a group of formula $-W-R^8$.

10 R^{3a} is hydrogen, C1-20-alkyl or a group of formula:



or NR^3R^{3a} is a group of formula



R^4 is hydrogen,

15 R^5 is hydrogen; nitro; halogen; azido; cyano; $-S-C1-4-alkyl$; $-SO-C1-4-alkyl$; $-SO_2-C1-4-alkyl$; $-SONH_2$; C1-20-alkyl unsubstituted or substituted by halogen; or C1-20-alkoxy unsubstituted or substituted by halogen,

R^6 is hydrogen, C1-20-alkyl or halogen,

R^7 is hydrogen, C1-20-alkyl or halogen,

20 W is C1-12-alkylene, $-NH-$ or $-NHC(=O)-$,

X is O, S or NH,

Y is O, S, $-CR^{12}R^{13}$, $-NR^{14}$ or $-C(=O)-$,

R^8 is aryl or heterocycle,

R^9 , R^{10} , R^{10a} and R^{11} are independently selected from hydrogen, C1-4-alkyl,

25 halogen, hydroxy or methoxycarbonyl,

or R^{10} and R^{10a} together form a C3-6-alkylene,

R^{12} is hydrogen, C1-4-alkyl, halogen or hydroxy,

R^{13} is hydrogen,

or CR¹²R¹³ is dioxolanyl,

R¹⁴ is aryl, heterocycle or a group of formula -V-R¹⁵,

V is C1-12-alkylene,

R¹⁵ is aryl or heterocycle,


5 m is 1 to 4,

n is 0 or 1,

and at least one of R⁵, R⁶ or R⁷ is different from hydrogen when R² is hydrogen,

R³ is H or 2,6-diisopropylphenyl, and R^{3a} is H.

- 10 2. A compound having the formula I or a pharmaceutically acceptable salt thereof or stereoisomeric forms thereof,




wherein

R¹ is hydrogen,


15 R² is hydrogen or C1-4-alkyl,

R³ is hydrogen; C1-6-alkyl unsubstituted or substituted by 1 to 5 substituents selected from halogen, hydroxy, alkoxy, alkoxycarbonyl or alkylamino; C5-7-cycloalkyl; (hydroxymethyl)cyclohexenyl; phenyl unsubstituted or substituted by 1 to 5 substituents selected from halogen, C1-4-alkyl, hydroxy, methoxy, nitro, methylsulfonyl, trifluoromethylthio or pyridinylalkyl; pyridinyl unsubstituted or substituted by methoxy; triazolyl; C1-4-alkoxy; or a group of formula -W-R⁸,
 20 R^{3a} is hydrogen, C1-4-alkyl or a group of formula



or NR³R^{3a} is piperidinyl unsubstituted or substituted by hydroxy;

25 thiomorpholinyl; thiazolidinyl unsubstituted or substituted by C1-4-alkoxycarbonyl; 2,5-dihydro-1H-pyrrol-1-yl; 1,4-dioxo-8-azaspiro[4.5]dec-8-yl; 4-oxooctahydro-1(2H)-quinolinyl; or a group of formula



30 R⁴ is hydrogen,

R⁵ is hydrogen; nitro; halogen; C1-4-alkyl, unsubstituted or substituted by halogen; or C1-4-alkoxy unsubstituted or substituted by halogen,

R⁶ is hydrogen, C1-6-alkyl or halogen,

R⁷ is hydrogen, methyl or halogen,

5 W is C1-4-alkylene unsubstituted or substituted by halogen, hydroxy, C1-4-alkyl or alkoxy; -NH-; or -NHC(=O)-,

R⁸ is phenyl unsubstituted or substituted by 1 to 5 substituents selected from halogen, C1-4-alkyl, hydroxy, methoxy, nitro, methylsulfonyl or trifluoromethylthio; furyl unsubstituted or substituted by methyl; pyrazolyl;

10 pyridinyl; morpholinyl; tetrahydrobenzazocinyl; piperidinyl unsubstituted or substituted by methyl; dihydroisochromenyl or dihydroimidazolyl,

R¹⁴ is pyridinyl; phenyl unsubstituted or substituted by halogen, hydroxy, C1-4-alkyl; or a group of formula -V-R¹⁵,

V is unsubstituted C1-4-alkylene,


15 R¹⁵ is phenyl or morpholinyl,

m is 1 to 4,

and at least one of R⁵, R⁶ or R⁷ is different from hydrogen when R² is hydrogen,

R³ is H or 2,6-diisopropylphenyl, and R^{3a} is H.

20 3. A compound having the formula I or a pharmaceutically acceptable salt thereof or stereoisomeric forms thereof,



wherein

R¹ is hydrogen,

25 R² is hydrogen, methyl or ethyl,

R³ is hydrogen, n-butyl, cycloheptyl, 2-fluoroethyl, 3-hydroxypropyl, 3-hydroxy-2,2-dimethylpropyl, 1-(hydroxymethyl)propyl, 3,3,3-trifluoro-2-hydroxypropyl, 3-ethoxypropyl, 2-ethoxy-2-oxoethyl, 3-(dimethylamino)propyl, 6-(hydroxymethyl)cyclohex-3-en-1-yl, 3-hydroxyphenyl, 3-fluorophenyl, 3-(2-pyridin-2-ylethyl)phenyl, 3,4-dimethylphenyl, 4-tert-butylphenyl, benzyl, 4-hydroxy-3-methoxybenzyl, 4-methylsulfonylbenzyl, 2-nitrobenzyl, 2-chloro-6-fluorobenzyl, 2-[(trifluoromethyl)thio]benzyl, 2-hydroxy-2-phenylethyl, 2-(3,4-

30

dimethoxyphenyl)ethyl, 2-(2-chlorophenyl)ethyl, 2-(4-methylphenyl)ethyl, (4-bromophenyl)amino, pyridin-3-yl, 6-methoxypyridin-3-yl, 4H-1,2,4-triazol-3-yl, pyridin-4-ylmethyl, (5-methyl-2-furyl)methyl, 3-(1H-pyrazol-1-yl)propyl, 2-morpholin-4-ylethyl, 2-((3,4,5,6-tetrahydro-1-benzazocin-1(2H)-yl)propyl, 2-(2-methylpiperidin-1-yl)ethyl, 3,4-dihydro-1H-isochromen-1-ylmethyl, methoxy, (4-pyridinylcarbonyl)amino or 4,5-dihydro-1H-imidazol-2-ylamino, R^{3a} is hydrogen, methyl or tetrahydrofuran-2-ylmethyl, or NR³R^{3a} 4-pyridin-2-ylpiperazin-1-yl, 4-(3-methylphenyl)piperazin-1-yl, 4-(4-hydroxyphenyl)piperazin-1-yl, 4-(2-phenylethyl)piperazin-1-yl, 4-(2-morpholin-4-ylethyl)piperazin-1-yl, 3-hydroxypiperidin-1-yl, thiomorpholin-4-yl, 4-methoxycarbonyl-1,3-thiazolidin-3-yl, 2,5-dihydro-1H-pyrrol-1-yl, 1,4-dioxo-8-azaspiro[4.5]dec-8-yl or 4-oxooctahydro-1(2H)-quinoliny, R⁴ is hydrogen, R⁵ is hydrogen, methyl, ethyl, trifluoromethyl, trifluoromethoxy, n-propyl, isopropyl, nitro or halogen, R⁶ is hydrogen, methyl or Cl, R⁷ is hydrogen, methyl, Br, F or Cl, and at least one of R⁵, R⁶ or R⁷ is different from hydrogen when R² is hydrogen, R³ is H or 2,6-diisopropylphenyl, and R^{3a} is H.

4. A compound according to any of the claims 1 to 3 wherein R² is hydrogen or methyl.
5. A compound according to any of the preceding claims wherein R³ is hydrogen.
6. A compound according to any of the preceding claims wherein R^{3a} is hydrogen.
7. A compound according to any of the preceding claims wherein R⁵ is halogen or trifluoromethyl.
8. A compound according to any of the preceding claims wherein R⁶ is hydrogen.
9. A compound according to any of the preceding claims wherein R⁷ is hydrogen, Br or F.
10. A compound according to any of the preceding claims wherein R² is C1-20-alkyl and the carbon atom to which R² is attached is in the "S"-configuration.

11. A compound selected from 2-(5-iodo-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5,7-dibromo-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-nitro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-methyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; (2R)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; (2S)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; 2-[2-oxo-5-(trifluoromethoxy)-2,3-dihydro-1H-indol-1-yl]acetamide; 2-(5-isopropyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-ethyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-fluoro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5,7-dimethyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-bromo-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(2-oxo-5-propyl-2,3-dihydro-1H-indol-1-yl)acetamide; 2-[2-oxo-5-(trifluoromethyl)-2,3-dihydro-1H-indol-1-yl]acetamide; 2-(5,6-dimethyl-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(7-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(6-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)butanamide; (+)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)butanamide; (-)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)butanamide; 2-(5-methyl-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; (+)-2-(5-methyl-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; (-)-2-(5-methyl-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; 2-(5-bromo-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; (-)-2-(5-bromo-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; (+)-2-(5-bromo-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide; 2-(5-chloro-7-fluoro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3-hydroxyphenyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3-fluorophenyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[3-(2-pyridin-2-ylethyl)phenyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[6-(hydroxymethyl)cyclohex-3-en-1-yl]acetamide; 5-chloro-1-[2-oxo-2-(4-pyridin-2-yl)piperazin-1-yl]ethyl-1,3-dihydro-2H-indol-2-one; 5-chloro-1-[2-[4-(3-methylphenyl)piperazin-1-yl]-2-oxoethyl]-1,3-dihydro-2H-indol-2-one; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(4-hydroxy-3-methoxybenzyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(pyridin-4-ylmethyl)-N-(tetrahydrofuran-2-ylmethyl)acetamide; 5-chloro-1-[2-(3-hydroxypiperidin-1-yl)-2-oxoethyl]-1,3-dihydro-2H-indol-2-one; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N'-isonicotinoylacetohydrazide; 5-chloro-1-(2-oxo-2-thiomorpholin-4-ylethyl)-1,3-dihydro-2H-indol-2-one; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(4H-1,2,4-triazol-3-yl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[4-(methylsulfonyl)benzyl]acetamide; 1-[(5-chloro-2-oxo-


2,3-dihydro-1H-indol-1-yl)acetyl]octahydroquinolin-4(1H)-one; N'-(4-bromophenyl)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetohydrazide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(6-methoxypyridin-3-yl)acetamide; N-butyl-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3-hydroxypropyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[3-(dimethylamino)propyl]acetamide; 5-chloro-1-[2-oxo-2-[4-(2-phenylethyl)piperazin-1-yl]ethyl]-1,3-dihydro-2H-indol-2-one; ethyl [(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetyl]amino}acetate; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3-ethoxypropyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(2-fluoroethyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-methoxy-N-methylacetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3,4-dimethylphenyl)acetamide; N-(4-tert-butylphenyl)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3-hydroxy-2,2-dimethylpropyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[1-(hydroxymethyl)propyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3,3,3-trifluoro-2-hydroxypropyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(2-hydroxy-2-phenylethyl)acetamide; 5-chloro-1-[2-[4-(4-hydroxyphenyl)piperazin-1-yl]-2-oxoethyl]-1,3-dihydro-2H-indol-2-one; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(pyridin-4-ylmethyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[(5-methyl-2-furyl)methyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[3-(1H-pyrazol-1-yl)propyl]acetamide; methyl 3-[(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetyl]-1,3-thiazolidine-4-carboxylate; 5-chloro-1-[2-(2,5-dihydro-1H-pyrrol-1-yl)-2-oxoethyl]-1,3-dihydro-2H-indol-2-one; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N'-(4,5-dihydro-1H-imidazol-2-yl)acetohydrazide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-(3,4-dimethoxyphenyl)ethyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-(2-chlorophenyl)ethyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-(4-methylphenyl)ethyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(2-morpholin-4-ylethyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-(3,4,5,6-tetrahydro-1-benzazocin-1(2H)-yl)propyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-(2-methylpiperidin-1-yl)ethyl]acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(2-nitrobenzyl)acetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-(3,4-dihydro-1H-isochromen-1-ylmethyl)acetamide; N-(2-chloro-6-fluorobenzyl)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide; N-benzyl-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-methylacetamide; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-[2-[(trifluoromethyl)thio]benzyl]acetamide; 5-chloro-1-[2-(1,4-dioxo-8-

azaspiro[4.5]dec-8-yl)-2-oxoethyl]-1,3-dihydro-2H-indol-2-one; 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-cycloheptylacetamide; 5-chloro-1-{2-[4-(2-morpholin-4-ylethyl)piperazin-1-yl]-2-oxoethyl}-1,3-dihydro-2H-indol-2-one; and 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)-N-pyridin-3-ylacetamide.

5

12. A compound selected from 2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetamide and (2S)-2-(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)propanamide.

13. A compound having the formula II or stereoisomeric forms thereof,



10

wherein


R¹ is hydrogen,

R² is hydrogen or C1-20-alkyl,


R³ is hydrogen, C1-20-alkyl, C4-8-cycloalkyl, C5-8-cycloalkenyl, aryl, aromatic or non aromatic heterocycle, C1-20-alkoxy, or a group of formula -W-R⁸.

15

R^{3a} is hydrogen, C1-20-alkyl or a group of formula:



or NR³R^{3a} is a group of formula



R⁴ is hydrogen,

20

R⁵ is hydrogen; nitro; halogen; azido; cyano; -S-C1-4-alkyl; -SO-C1-4-alkyl; -SO₂-C1-4-alkyl; -SONH₂; C1-20-alkyl unsubstituted or substituted by halogen; or C1-20-alkoxy unsubstituted or substituted by halogen,

R⁶ is hydrogen, C1-20-alkyl or halogen,

R⁷ is hydrogen, C1-20-alkyl or halogen,

25

W is C1-12-alkylene, -NH- or -NHC(=O)-,


X is O, S or NH,

Y is O, S, -CR¹²R¹³-, -NR¹⁴- or -C(=O)-,

R⁸ is aryl or heterocycle,

R^9 , R^{10} , R^{10a} and R^{11} are independently selected from hydrogen, C1-4-alkyl, halogen, hydroxy or methoxycarbonyl,
 or R^{10} and R^{10a} together form a C3-6-alkylene,
 R^{12} is hydrogen, C1-4-alkyl, halogen or hydroxy,
 R^{13} is hydrogen,
 or $CR^{12}R^{13}$ is dioxolanyl,
 R^{14} is aryl, heterocycle or a group of formula $-V-R^{15}$,
 V is C1-12-alkylene,
 R^{15} is aryl or heterocycle,
 m is 1 to 4,
 n is 0 or 1,
 and at least one of R^5 , R^6 or R^7 is different from hydrogen when R^2 is hydrogen,
 R^3 is H or 2,6-diisopropylphenyl, and R^{3a} is H.

14. A compound having the formula III or stereoisomeric forms thereof,



wherein

R^4 is hydrogen,

R^5 is hydrogen; nitro; azido; cyano; $-S-C1-4-alkyl$; $-SO-C1-4-alkyl$;

$-SO_2-C1-4-alkyl$; $-SONH_2$; C1-20-alkyl unsubstituted or substituted by halogen;


or C1-20-alkoxy unsubstituted or substituted by halogen,

R^6 is hydrogen, C1-20-alkyl or halogen,

R^7 is hydrogen, C1-20-alkyl or halogen,

and at least one of R^5 , R^6 or R^7 is different from hydrogen.

15. A compound having the formula VI or stereoisomeric forms thereof,



wherein

R^1 is hydrogen,


R^2 is hydrogen or C1-20-alkyl,

R^3 is hydrogen, C1-20-alkyl, C4-8-cycloalkyl, C5-8-cycloalkenyl, aryl, aromatic or non aromatic heterocycle, C1-20-alkoxy, or a group of formula $-W-R^8$.

R^{3a} is hydrogen, C1-20-alkyl or a group of formula:



or NR^3R^{3a} is a group of formula:



R^4 is hydrogen,

R^5 is hydrogen; halogen; azido; cyano; $-S-C1-4-alkyl$; $-SO-C1-4-alkyl$;

10 $-SO_2-C1-4-alkyl$; $-SONH_2$; or C1-20-alkyl unsubstituted or substituted by halogen,

R^6 is hydrogen, C1-20-alkyl or halogen,

R^7 is hydrogen, C2-20-alkyl or halogen,

W is C1-12-alkylene, $-NH-$ or $-NHC(=O)-$,

15 X is O, S or NH,

Y is O, S, $-CR^{12}R^{13}$ -, $-NR^{14}$ - or $-C(=O)-$,

R^8 is aryl or heterocycle,

R^9 , R^{10} , R^{10a} and R^{11} are independently selected from hydrogen, C1-4-alkyl, halogen, hydroxy or methoxycarbonyl,

20 or R^{10} and R^{10a} together form a C3-6-alkylene,

R^{12} is hydrogen, C1-4-alkyl, halogen or hydroxy,

R^{13} is hydrogen,

or $CR^{12}R^{13}$ is dioxolanyl,

R^{14} is aryl, heterocycle or a group of formula $-V-R^{15}$,

25 V is C1-12-alkylene,

R^{15} is aryl or heterocycle,


m is 1 to 4,

n is 0 or 1,

and at least one of R^5 , R^6 or R^7 is different from hydrogen when R^2 is hydrogen,

30 R^3 is H or 2,6-diisopropylphenyl, and R^{3a} is H.

16. A compound having the formula IX or stereoisomeric forms thereof,



wherein

R¹ is hydrogen,

R² is hydrogen or C1-20-alkyl,

5 R⁴ is hydrogen,

R⁵ is hydrogen; nitro; azido; cyano; -S-C1-4-alkyl; -SO-C1-4-alkyl;
-SO₂-C1-4-alkyl; -SONH₂; halogen; C1-20-alkyl unsubstituted or substituted by
halogen; or C1-20-alkoxy unsubstituted or substituted by halogen,

R⁶ is hydrogen, C1-20-alkyl or halogen,

10 R⁷ is hydrogen, C1-20-alkyl or halogen,

and at least one of R⁵, R⁶ or R⁷ is different from hydrogen when R² is hydrogen,

R³ is H or 2,6-diisopropylphenyl, and R^{3a} is H.

17. A compound which is selected from the group consisting of:


- 15 2-(5'-methyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide;
2-[2'-oxo-5'-[(trifluoromethyl)oxy]spiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-
yl]acetamide;
2-[5'-(1-methylethyl)-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl]acetamide;
2-(5'-ethyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide;
20 2-(5'-fluoro-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide;
2-(5',7'-dimethyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide; 2-(2'-
oxo-5'-propylspiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide;
2-[2'-oxo-5'-(trifluoromethyl)spiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl]acetamide;
2-(5',6'-dimethyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide;
25 5'-methylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
5'-[(trifluoromethyl)oxy]spiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
5'-(1-methylethyl)spiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
5'-ethylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
5'-fluorospiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
30 5',7'-dimethylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
5'-propylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
5'-(trifluoromethyl)spiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
5',6'-dimethylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;

- 2-(5-chloro-1H-indol-1-yl)propanamide;
2-(7-chloro-1H-indol-1-yl)acetamide;
2-(6-chloro-1H-indol-1-yl)acetamide;
2-(5-chloro-1H-indol-1-yl)butanamide;
5 2-(5-methyl-1H-indol-1-yl)propanamide;
2-(5-bromo-1H-indol-1-yl)propanamide;
2-(7-fluoro-1H-indol-1-yl)acetamide;
2-(5-bromo-1H-indol-1-yl)acetamide;
2-(5-fluoro-1H-indol-1-yl)acetamide;
10 2-(5-chloro-1H-indol-1-yl)acetamide;
(5-chloro-2-oxo-2,3-dihydro-1H-indol-1-yl)acetic acid.
18. A pharmaceutical composition comprising an effective amount of a compound
according to any of claims 1 to 12 in combination with a pharmaceutically
15 acceptable diluent or carrier.
19. A method for treating epilepsy, epileptogenesis, seizure disorders, convulsions,
Parkinson's disease, dyskinesia induced by dopamine replacement therapy, tardive
dyskinesia induced by administration of neuroleptic drugs, Huntington Chorea,
20 and other neurological disorders including bipolar disorders, mania, depression,
anxiety, attention deficit hyperactivity disorder (ADHD), migraine, trigeminal and
other neuralgia, chronic pain, neuropathic pain, cerebral ischemia, cardiac
arrhythmia, myotonia, cocaine abuse, stroke, myoclonus, tremor, essential tremor,
simple or complex tics, Tourette syndrome, restless leg syndrome and other
25 movement disorders, neonatal cerebral haemorrhage, amyotrophic lateral
sclerosis, spasticity and degenerative diseases, bronchial asthma, asthmatic status
and allergic bronchitis, asthmatic syndrome, bronchial hyperreactivity and
bronchospastic syndromes as well as allergic and vasomotor rhinitis and
rhinoconjunctivitis, in a mammal in need of such treatment, comprising
30 administering a therapeutic dose of at least one compound according to any of
claims 1-12 or a pharmaceutical composition according to claim 18.
20. A compound according to any of claims 1-12 or a pharmaceutical composition
according to claim 18 for use as a medicament.
35
21. Use of a compound according to any of claims 1-12 or of a pharmaceutical
composition according to claim 18 for the manufacture of a medicament for the

5 treatment of epilepsy, epileptogenesis, seizure disorders, convulsions, Parkinson's
disease, dyskinesia induced by dopamine replacement therapy, tardive dyskinesia
induced by administration of neuroleptic drugs, Huntington Chorea, and other
neurological disorders including bipolar disorders, mania, depression, anxiety,
10 attention deficit hyperactivity disorder (ADHD), migraine, trigeminal and other
neuralgia, chronic pain, neuropathic pain, cerebral ischemia, cardiac arrhythmia,
myotonia, cocaine abuse, stroke, myoclonus, tremor, essential tremor, simple or
complex tics, Tourette syndrome, restless leg syndrome and other movement
15 disorders, neonatal cerebral haemorrhage, amyotrophic lateral sclerosis, spasticity
and degenerative diseases, bronchial asthma, asthmatic status and allergic
20 bronchitis, asthmatic syndrome, bronchial hyperreactivity and bronchospastic
syndromes as well as allergic and vasomotor rhinitis and rhinoconjunctivitis.

AMENDED CLAIMS

[received by the International Bureau on 26 August 2004 (26.08.2004);
original claim 16 amended; remaining claims unchanged (1 page)]



wherein

R¹ is hydrogen,

R² is hydrogen or C1-20-alkyl,

5 R⁴ is hydrogen,

R⁵ is hydrogen; nitro; azido; cyano; -S-C1-4-alkyl; -SO-C1-4-alkyl;
-SO₂-C1-4-alkyl; -SONH₂; halogen; C1-20-alkyl unsubstituted or substituted by
halogen; or C1-20-alkoxy unsubstituted or substituted by halogen,

R⁶ is hydrogen or C1-20-alkyl,

10 R⁷ is hydrogen, C1-20-alkyl or halogen,

and at least one of R⁵, R⁶ or R⁷ is different from hydrogen when R² is hydrogen,
R³ is H or 2,6-diisopropylphenyl, and R^{3a} is H.

17. A compound which is selected from the group consisting of:

- 15 2-(5'-methyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide;
2-[2'-oxo-5'-((trifluoromethyl)oxy)spiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-
yl]acetamide;
2-[5'-(1-methylethyl)-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl]acetamide;
2-(5'-ethyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide;
20 2-(5'-fluoro-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide;
2-(5',7'-dimethyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide; 2-(2'-
oxo-5'-propylspiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide;
2-[2'-oxo-5'-(trifluoromethyl)spiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl]acetamide;
2-(5',6'-dimethyl-2'-oxospiro[1,3-dithiolane-2,3'-indol]-1'(2'H)-yl)acetamide;
25 5'-methylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
5'-[(trifluoromethyl)oxy]spiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
5'-(1-methylethyl)spiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
5'-ethylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
5'-fluorospiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
30 5',7'-dimethylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
5'-propylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
5'-(trifluoromethyl)spiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;
5',6'-dimethylspiro[1,3-dithiolane-2,3'-indol]-2'(1'H)-one;

INTERNATIONAL SEARCH REPORT

 II
 national Application No
 PCT/EP2004/002691

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D209/34 A61K31/405 C07D209/54 C07D209/08 A61P25/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, BEILSTEIN Data, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL PATENTS INDEX, DOCUMENTATION ABSTRACTS JOURNAL Section Ch, Week 9627 Derwent Publications Ltd., London, GB; Class B02, AN 96-266522 XP002241135 & SU 841 264 A (A MED PHARMACOLOGY RES INST) 10 November 1995 (1995-11-10) cited in the application abstract	1-21
A	US 4 624 963 A (KRUSE HANSJOERG ET AL) 25 November 1986 (1986-11-25) page 1, column 1, line 11 - page 1, column 2, line 7 ----- -/--	1-21

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

29 June 2004

Date of mailing of the international search report

08/07/2004

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Usueli, A

INTERNATIONAL SEARCH REPORT

In ☐ International Application No
PCT/EP2004/002691

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94/29272 A (ASTRA AB ;BOAR BERNARD ROBIN (GB); SHEA DENNIS MARK O (GB); TOMLIN) 22 December 1994 (1994-12-22) page 2, line 7 - page 3, line 16 -----	1-21
X	EP 0 445 781 A (STERLING DRUG INC) 11 September 1991 (1991-09-11) Preparation 3D page 11, line 18 - page 11, line 18 -----	15
X	DATABASE CROSSFIRE BEILSTEIN 'Online! Beilstein Institut zur Förderung der Chemischen Wissenschaften, Frankfurt am Main, DE; XP002241130 Database accession no. BRN:6400176 abstract & NILSSON, I. ET AL.: ACTA CHEM. SCAND. SER. B, vol. 39, no. 7, 1985, pages 531-548, -----	15
X	DATABASE CROSSFIRE BEILSTEIN 'Online! Beilstein Institut zur Förderung der Chemischen Wissenschaften, Frankfurt am Main, DE; XP002241131 Database accession no. BRN:4296865 abstract & BELL, M. ET AL.: J.MED.CHEM., vol. 34, no. 3, 1991, pages 1099-1110, -----	15
X	HOTZEL C ET AL: "Design, synthesis, DNA-binding and cytotoxicity evaluation of new potential combilexines" EUROPEAN JOURNAL OF MEDICINAL CHEMISTRY, EDITIONS SCIENTIFIQUE ELSEVIER, PARIS, FR, vol. 37, no. 5, May 2002 (2002-05), pages 367-378, XP004372953 ISSN: 0223-5234 compound 16 figure 2 -----	15
X	WO 98/01428 A (DU PONT MERCK PHARMA) 15 January 1998 (1998-01-15) page 73, line 33 - page 73, line 34; example 202 -----	15
X	EP 0 022 317 A (FUJISAWA PHARMACEUTICAL CO) 14 January 1981 (1981-01-14) page 21, line 11; compound 9 -----	16
A	PATENT ABSTRACTS OF JAPAN vol. 0133, no. 08 (C-617), 14 July 1989 (1989-07-14) & JP 1 095766 A (TAMANOISU KK), 13 April 1989 (1989-04-13) abstract -----	1

INTERNATIONAL SEARCH REPORT

national application No.
PCT/EP2004/002691

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: —
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 19 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/EP2004/002691

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
SU 841264	A	10-11-1995	SU 841264 A1	10-11-1995
US 4624963	A	25-11-1986	DE 3413572 A1	24-10-1985
			AU 4106085 A	17-10-1985
			EP 0158278 A2	16-10-1985
			JP 60228459 A	13-11-1985
			ZA 8502638 A	27-11-1985
WO 9429272	A	22-12-1994	AU 7010894 A	03-01-1995
			CA 2164119 A1	22-12-1994
			EP 0703901 A1	03-04-1996
			FI 956074 A	18-12-1995
			JP 8511515 T	03-12-1996
			NO 955074 A	07-02-1996
			WO 9429272 A1	22-12-1994
EP 0445781	A	11-09-1991	AT 133663 T	15-02-1996
			AU 643757 B2	25-11-1993
			AU 7266991 A	12-09-1991
			CA 2036307 A1	09-09-1991
			DE 69116705 D1	14-03-1996
			DE 69116705 T2	26-09-1996
			DK 445781 T3	19-02-1996
			EP 0445781 A1	11-09-1991
			ES 2082872 T3	01-04-1996
			FI 911131 A ,B,	09-09-1991
			GR 3019058 T3	31-05-1996
			HU 56492 A2	30-09-1991
			IE 910700 A1	11-09-1991
			IL 97426 A	31-03-1996
			JP 3157179 B2	16-04-2001
			JP 4235164 A	24-08-1992
			KR 183396 B1	01-05-1999
			MX 24839 A ,B	01-10-1993
			NO 910868 A ,B,	09-09-1991
			NZ 237309 A	25-02-1993
			PH 27022 A	01-02-1993
			PT 96938 A ,B	31-10-1991
			RU 2073670 C1	20-02-1997
			US 5013837 A	07-05-1991
WO 9801428	A	15-01-1998	AU 3645697 A	02-02-1998
			CA 2259573 A1	15-01-1998
			EP 0960102 A1	01-12-1999
			JP 2002514162 T	14-05-2002
			NZ 333696 A	23-06-2000
			WO 9801428 A1	15-01-1998
EP 0022317	A	14-01-1981	JP 55167282 A	26-12-1980
			AT 4713 T	15-10-1983
			DE 3064910 D1	27-10-1983
			EP 0022317 A1	14-01-1981
			JP 1014223 B	10-03-1989
			JP 1531352 C	24-11-1989
			JP 56097268 A	05-08-1981
			US 4370340 A	25-01-1983
			US 4438126 A	20-03-1984

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/EP2004/002691

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 1095766	A	13-04-1989	NONE